



Western Michigan University
ScholarWorks at WMU

International Conference on African
Development Archives

Center for African Development Policy
Research

8-2007

Maternal Education and Child Nutrition: Evidence from the 2000 and 2005 Ethiopian Demographic and Health Surveys

Alemayehu Azeze Ambel
Western Michigan University, alemayehu.ambel@wmich.edu

Follow this and additional works at: https://scholarworks.wmich.edu/africancenter_icad_archive



Part of the African Studies Commons, and the Economics Commons

WMU ScholarWorks Citation

Ambel, Alemayehu Azeze, "Maternal Education and Child Nutrition: Evidence from the 2000 and 2005 Ethiopian Demographic and Health Surveys" (2007). *International Conference on African Development Archives*. 98.

https://scholarworks.wmich.edu/africancenter_icad_archive/98

This Paper is brought to you for free and open access by the Center for African Development Policy Research at ScholarWorks at WMU. It has been accepted for inclusion in International Conference on African Development Archives by an authorized administrator of ScholarWorks at WMU. For more information, please contact wmu-scholarworks@wmich.edu.



**Maternal Education and Child Nutrition: Evidence from the 2000 and 2005
Ethiopian Demographic and Health Surveys**

DRAFT

Alemayehu Azeze Ambel
alemayehu.ambel@wmich.edu
Department of Economics
Western Michigan University

Abstract

I used the 2000 and 2005 Ethiopian Demographic and Health Surveys to analyze the effect of maternal education and its pathways on chronic and acute malnutrition in Ethiopia. The pathways examined in this study are socioeconomic status, maternal health-seeking behavior, maternal knowledge of health and family planning and reproductive behavior. I find that maternal education works through all except health-seeking behavior. I also find that maternal education and its pathways are more relevant and robust in explaining chronic than acute malnutrition. Socioeconomic status is the most important factor linking maternal education and child nutritional status. Although girls' education is a high policy priority, it may take time before its direct and indirect impacts substantially improve child health outcomes. Faster results would require direct interventions on key elements of socioeconomic status

Keywords: Maternal Education, Nutrition, Children, Ethiopia
JEL: I12; J13

1. Introduction

The positive and strong relationship between maternal education and child health outcomes is widely documented and largely undisputed (Frongillo, de Onis and Hanson, 1997; Variyam et al, 1999; Alderman et al., 2000; Smith and Haddad, 2000). A large body of literature documents that maternal education works through a number of pathway variables that directly affects child health outcomes. The list includes a number of maternal, household and community characteristics such as socioeconomic status, geographic residence, nutritional and health knowledge, autonomy, health-seeking and reproductive behavior (Desai and Alva, 1998; Glewwe, 1999; Webb and Block, 2004; Frost, Forste and Haas, 2005).

Therefore, the impact of education on child health is greatly attenuated when selected mediating factors are included in the model. However, there is a broad disagreement on the role of the various linkages through which the impacts of maternal education on child health outcome are transmitted. Moreover, it is noted that the impact of maternal education could be different to different markers of child nutritional status (Webb and Block, 2004).

This study models selected pathways linking maternal education and child nutritional status in Ethiopia using the 2000 and 2005 Ethiopian Demographic and Health Survey (EDHS). The study empirically investigates how maternal education and its various pathways affect chronic (height for age) and acute (weight for height) malnutrition in children younger than five years. The study addresses the following questions: (1) Does maternal education work through selected pathways such as socio-economic status, maternal health-seeking behavior, and maternal knowledge of family planning and health? (2) How do maternal education and its pathways perform in models of different types of child malnutrition?

The study is motivated by some observations from recent developments on child nutritional status in Ethiopia. First, the 2005 EDHS shows significant improvement in child nutritional status when compared to the results of preceding survey conducted in 2000. Second, the changes have been different to different measures of child malnutrition. Chronic malnutrition declined while acute malnutrition remained the same

on average (Central Statistical Authority and ORC Macro, 2001&2006). The availability of comparable survey data would provide an opportunity to investigate the performance of the determinants of child nutritional status over time.

Fourth, malnutrition is a leading cause of child death in developing countries (Black, Morris and Bryce, 2003) and reducing child mortality is among the major priorities included in the Millennium Development Goals (MDGs). The fact that the prevalence of child malnutrition and infant mortality in Ethiopia is among the highest in developing regions, the issue is of national and international concern.

Previous studies on child nutritional status in Ethiopia focused on identifying the determinants of chronic malnutrition from a one time survey (e.g. Girma and Genebo, 2002) or without due emphasis on the impact of education in various contexts (e.g. Christaensen and Alderman, 2004). My study expands the discussion in two ways. First, the study analyzes the effect of maternal education on child nutritional status considering a more comprehensive array of linkages. Second, the study considers both chronic and acute malnutrition.

The rest of the paper proceeds as follows: Section 2 provides a background on the prevalence and recent changes in child nutritional status in Ethiopia. Section 3 is on the multivariate analysis including the empirical framework and the data. Section 4 discusses the results. Finally, concluding remarks are provided in Section 5.

2. Background: Child Nutritional Status in Ethiopia

The conventional measures of child anthropometrics show that Ethiopia ranks among those countries in sub-Saharan Africa with the high prevalence of child malnutrition. In 2003, 52% of children were suffering from chronic malnutrition (stunting), 11% from acute malnutrition (wasting) and 47% from underweight. During the same period, the average prevalence of stunting, wasting and underweight for African countries were 39%, 9% and 29% respectively.¹ A recently completed survey in Ethiopia, the 2005 EDHS, shows a similar profile of under-five malnutrition (Table 1).

Another important feature of child nutritional status in Ethiopia is that the prevalence can be distinguished by selected background characteristics. Table 1 presents

the performance of child nutritional status by maternal education and place of residence in 2000 and 2005. It shows that the prevalence of malnutrition among children whose mother has some education is lower than those whose mother has no at least primary education. The percent of children malnourished consistently declines, as the highest level of education attained by the mother increases from no education to primary education, and then to secondary and higher education. The trend is consistent across different indicators of child malnutrition and survey years.

<Table 1>

Table 1 also shows that the prevalence of malnutrition in general is lower in 2005 than in 2000. The percentage declines over the period 2000-2005 show that the reductions in child malnutrition (for stunting and underweight) is the highest for the highest level of maternal education (which is “secondary and higher education”) and the lowest for the lowest level of education (which is “no education”). However, there is no consistent decline for wasting. This could be due to the fact that stock variables such as education and place of residence are stock variables and better explain chronic outcomes such as stunting than acute fluctuations in nutritional status.

As expected, the urban advantage presented in Table 1 is unambiguous when the performance of child nutritional status is disaggregated by place of residence without controlling for other factors. In addition, the comparison between the 2000 and 2005 survey results shows that the reductions in children stunting and underweight are larger in urban than in rural areas. However, a number of studies find that any conclusion based on a simple bivariate relationship would be misleading because the “advantage” often disappears when other important variables are included (Fotso, 2006).

The kernel density plots in Figure 1 and Figure 2 corroborate the results in Table 4.1. Figure 4.1 shows that the distributions can be differentiated by maternal education in national, rural and urban samples. In each panel, the dashed lines are to the right of the solid lines showing expected differences in height for age z-scores (HAZ) and weight-for height z-scores (WHZ) of children by maternal education. The impact of maternal education is larger in HAZ and than in WHZ (compare columns: Panels A1, B1 & C1

Vs Panels A2, B2 & C2). It is also larger in rural than in urban areas (compare rows Panel B1&B2 Vs Panels C1&C2).

Similarly, Figure 2 shows kernel density plots of HAZ and WHZ for national (all), rural and urban children by survey year. Each panel comprises two plots for each survey year. The objective of the plots in each panel is to show if there were changes in the distribution between the two surveys. Panel D1 and D2 are for the national sample; Panel E1 and E2 are for the rural sample; and Panel F1 and F2 are for the urban sample. In all the three cases, the densities in 2005 are to the right of that of the 2000 implying improvements in child nutrition in 2005.

<Figure 1>
<Figure 2>

The significance of the differences of child nutritional status by background characteristics and time presented above (Figures 1 & 2 and Table 1) is checked by a Kolmogorov-Smirnov (KS) test of equality between the two empirical distributions is carried out. In this regard, the three null hypotheses that are being tested include (1) $H_0: F_{\text{some education}}(z) = F_{\text{no education}}(z) \forall z$, (2) $H_0: F_{\text{urban}}(z) = F_{\text{rural}}(z) \forall z$, and (3) $H_0: F_{2005}(z) = F_{2000}(z) \forall z$. The KS test is based on the largest absolute gap between the cumulative distributions of F_1 and F_0 where there are m observations for distribution 1 and n

observations for distribution 0, i.e., $F_0(z) = \frac{1}{n} \sum_{i=1}^n \begin{cases} 1 & y_i \leq z \\ 0 & \text{otherwise} \end{cases}$ and $F_1(z) = \frac{1}{m} \sum_{i=1}^m \begin{cases} 1 & y_i \leq z \\ 0 & \text{otherwise} \end{cases}$,

where z is an indicator of child nutritional status, including HAZ, WHZ, and WAZ. Then, the test statistic is obtained from the supremum of the absolute values of the differences of the two empirical cumulative distribution functions, i.e.,

$$D = \max_z |F_0(z) - F_1(z)|.$$

<Table 2>

Table 2 presents results of the KS test of equality of distributions for wasting², stunting and underweight for national, rural and urban samples. The p-values show that, in all the three cases, the null hypothesis that the two distributions are the same is rejected at less

than 1% level of significance. Coupled with the density plots in Figure 2, the result indicates significant improvement in child nutritional status.

It is important to note that the reductions in stunting and underweight during the 2000–2005 period are not only statistically significant but also have important economic implications. In 2005 about 13 million (17%) of the total Ethiopian population of 77.4 million were between age 0-4. Therefore, the decline in stunting by 5 percentage points, i.e., from 51.5% in 2000 to 46.5% in 2005, implies that there were 653,000 fewer stunted children in 2005 than those that there would have been if the percent of stunting remained the same as in 2000.

Similarly, the decline in underweight by 8.8 percentage points (from 47.2% in 2000 to 38.4 % in 2005) would mean that 1.15 million fewer underweight children than those that there would have been if the percent of underweight remained the same as in 2000. Therefore, the question that remain are: what are the relevant factors, and to what extent maternal education and its pathways explain child malnutrition in Ethiopia?

3. Multivariate Analysis

3.1 The Empirical Framework

The standard procedure of identifying the determinants of child health outcomes involves maximizing the household's utility function subject to the biological or anthropometric production function and other constraints (Pitt and Rosenzweig 1985; Behrman and Deolalikar, 1988; Thomas, Lavy and Strauss, 1996; Webb and Block, 2004). Equation (1) presents the household's utility maximization problem, which is function of H_i (health status), F_i (food intake), L_i (leisure), G_i (consumption of other goods). Health of household members and food intake enter directly into the utility function because health is good in itself and food is taken for reasons other than nutritional value. The utility function may also be conditioned by observable individual characteristics (X_i), household characteristics (X_h), community characteristics (X_c) and unobserved heterogeneity of preferences (ψ_i).

$$\max_{H_i, F_i, L_i, G_i} U = U(H_i, F_i, L_i, G_i; X_i, X_h, X_c, \psi_i) \quad U' > 0, U'' < 0 \quad (1)$$

The household maximizes the utility function subject to a budget constraint and a biological health production function given by,

$$H_i = H(F_i, M_i, X_i, X_h, X_c, \eta_i) \quad (2)$$

where H_i is nutritional status as measured in anthropometrics outcomes (e.g. height or weight), M_i are non-food health inputs, η_i is unobserved individual health endowments, and all other variables are as defined earlier. Then, the maximization problem leads to a reduced form demand function for nutritional status:

$$H_i = h(X_i, X_h, X_c, v_i) \quad (3)$$

where v_i is unobserved nutritional outcome. Equation 3 provides a benchmark specification for empirical analysis.³ Equation 3 basically specifies nutritional status as a function of individual, household, and community characteristics. An important limitation of this approach is that it does not allow inferring structural coefficients. However, the reduced form equation is still informative about the effects on nutrition of changes in the explanatory variables thereof.

There exists tremendous variation in the specification of the empirical model of child nutritional status. Variants of the empirical models derived from Equation 3 often emanate from the choice of the dependent variable as well as the definition and measurement of individual, household and community characteristics. The availability of data also dictates the empirical specification. Equation 3 can be rearranged to specify an empirical model that distinguishes maternal education, pathway variables, and other control variables.

3.2 Incorporating pathway variables

Four key pathways are considered. These are socioeconomic status, health-seeking behavior, knowledge of health and family planning, and reproductive behavior.

Socioeconomic status

Maternal education has a clear connection with the various key elements of socioeconomic status including high-income job, possession of assets, better health and sanitary conditions, to mention but a few. The empirical evidence demonstrates the

existence of strong positive relationship between socioeconomic status and child health outcomes. Therefore, socioeconomic status stands out to be an important mediating factor between maternal education and child health.

Health-seeking behavior

Education can also influence health care utilization and reproductive health behavior. As Pongou, Ezzati and Salomon (2006) note, in some traditional societies, education would provide the mother with the capacity to break with some traditional practices and taboos. Education promotes modern attitudes and hence mothers with higher levels of education are more likely to seek healthcare services from health centers and health professionals. Educated mothers are also more likely to accept and use family planning methods including contraceptives.

Knowledge of family planning and health

Education enhances mother's knowledge of health, which is an important predictor of child health outcome (Glewwe, 1999; Webb and Block, 2004). Health knowledge can directly be acquired from formal education. Education can also facilitate the mother's ability to understand the causation and prevention methods of illness. It also enhances her knowledge of nutrition and family planning. However, Frost, Forste and Haas (2005) review that the available empirical evidence on the relationship between maternal knowledge of and child health is inconclusive.

Reproductive behavior

Reproductive behavior is another important link through which education influences child health outcome. In general, educated women have more control over their reproductive behavior and make conscious decisions, for example, on the number of births and intervals. Reproductive behavior is proxied by mother's age and selected and child demographic characteristics. Relevant child characteristics include age, sex, birth order and preceding birth interval.⁴

Vast evidence shows that the risk of child malnutrition increases with age in developing countries. Webb and Block (2004, p.812) find that HAZ and WHZ decline

with age though with a positive second derivative. An explanation for this relationship is the nutritional value of breastfeeding that protects young children from the risk of stunting or wasting at early age (e.g. Pongou, Ezzati and Salomon, 2006) and potentially, shortage of supplemental food in later months. In addition, some measures of malnutrition such as stunting are results of cumulative process of inadequate dietary intake and illness. Therefore, younger children are at lower risk (Webb and Block, 2004).

The rationale for including gender in the model of child nutrition is to capture the presence of male-bias in intrahousehold allocation of resources (Behrman, 1997).⁵ However, the empirical evidence to support this hypothesis remains scarce. Based on a review of 306 child nutrition surveys conducted since 1985 in a number of developing countries, Marcoux (2002) finds no sex differences in 227 surveys. In fact, the evidence from Africa and some other developing countries in Asia and Latin America shows that, when significant differences exist, boys are more likely to be malnourished than girls., when significant differences exist, boys are more likely to be malnourished than girls.

Birth order measures parity while birth interval captures the care and support that have been made available to the child. The empirical evidence on parity is mixed. For example, in India, Jeyaseelan and Lakshman (1997) find that malnutrition is higher among children of higher birth order (5+). On the other hand, In Ethiopia, Girma and Genebo (2002) find that the risk of stunting is higher among first births. However, it is common to find a result that supports the claim that the risk of malnutrition declines with birth interval (e.g. Pongou, Ezzati and Salomon, 2006).

Finally, place of residence and geographic regions are included as control variables in most specifications. It should be noted however that these controls are also influenced by maternal education. Education increases mobility and creates more opportunities in urban than rural areas. Desai and Alva (1998) find that in addition to socioeconomic factors, geographical controls are important links through which the impact of maternal education on child health outcome is mediated.

3.4.3 Data and Measurement of Variables

Data

The descriptions and analyses of this study are based on the two waves of the Ethiopian Demographic and Health Survey (EDHS) available at the time of writing (Central Statistical Agency and ORC Macro, 2001 & 2006). The first survey was completed in 2000 and the second survey was completed in 2005.

The EDHS sample is stratified, clustered and collected in two-stage probabilistic sampling technique based on the list of enumeration areas of the 1994 Population and Housing Census of Ethiopia. Therefore, the description and analysis undertaken in this study take into account the nature of the data. Accordingly, the sample weight, sample strata and primary sampling units are included.

At the first sampling stage in the 2000 survey, 539 (138 urban and 401 rural) clusters were selected. In the 2005 survey, 540 (145 urban and 395 rural) clusters were selected. The second stage consisted of the selection of a representative sample of households and women aged 15-49 years old in each household. Accordingly, in the 2000 survey, 15,367 women from 14,072 households were selected. In the 2005 survey, 14,070 women from 14,500 households were selected.

In both surveys, women were asked questions on their children especially for children younger than 5 years old and anthropometrics measurements (height and weight) were taken. In the 2000 and 2005 surveys, the total number of children measured and whose mothers were also interviewed were 9,774 and 4,296 respectively.

Measurement of variables

Dependent Variable: Child Nutritional Status

Long-term or chronic malnutrition is measured by height for age (HAZ) while short-term or acute malnutrition is measured by weight for height (WHZ). A child is said stunted if HAZ score is less than $-2SD$ and wasted if WHZ score is less than $-2SD$. Therefore, the dependent variable is a dichotomous variable that takes one if the child is stunted or wasted, and zero otherwise.

Explanatory Variables

The primary variables of interest are *maternal education* and *pathway variables*. The models are also controlled for geography (place of residence and regions) and survey year. The variables are measured as follows.

The DHS data compile *maternal education* in two different forms: single years and highest level of education. For ease of interpretation, six categories are considered following the DHS classification. These are: no education, incomplete primary education, complete primary education, incomplete secondary education, complete secondary education, and higher education. The corresponding values from the smallest to the highest education category range from 0-5.

Socioeconomic status is measured differently in different studies. Frost, Forste and Haas (2005) construct two index variables from selected household assets and dwelling characteristics. However, for this study the DHS wealth index is used because in addition to a number of household assets and dwelling characteristics, it considers the household's demographic structure.⁶ Assets and amenities included in the DHS wealth index range from the possession of items (e.g. bicycles, cars, radios, sofas, and televisions); dwelling characteristics such as type of flooring material or the level of overcrowding; household facilities such as source of drinking water, type of toilet facility, and type of cooking fuel; and other characteristics related to wealth status.

Mother's health-seeking behavior is an index variable constructed from utilization of selected preventive health care services. It is constructed by principal component analysis from four related variables included in the EDHS (Table 3). These are, (1) received prenatal services from a health professional or a trained birth attendant; (2) delivered a baby at a health center (hospital, clinic, others), (3) have used contraceptive, and (4) received tetanus injection before birth.

Similarly, *maternal knowledge* of family planning and health is an index variable constructed from selected variables available in the 2000 and 2005 EDHS. Health knowledge is measured by knowledge of oral rehydration therapy, i.e., if the woman heard of or used oral rehydration therapy. Family planning knowledge is measured by knowledge of ovulatory cycle, i.e. if the woman knows when in ovulatory cycle she can

get pregnant. Additional factors included in the knowledge index are proxies of family planning information from radio, TV, newspaper and frequencies of reading newspaper, listening to radio and watching TV (Table 3).

<Table 3>

Reproductive behavior is proxied by maternal age and selected child characteristics such age, sex of child, birth order, and birth interval. Child age is in months and maternal age is in years. Both are in logs. The remaining, namely, sex of child sex, birth order and preceding birth intervals are dummy variables.

4. Results

Table 4 presents descriptive statistics of the primary variables included in the regressions excluding control dummy variables for place of residence and geographic regions. As discussed earlier in Section 2, the first four variables in Table 4 show improvement in child nutritional status over the period 2000- 2005. For example, percent stunted declined from 51% to 46%; and HAZ increased from -2.06 to -1.77 . Similarly, although percent wasted remained at about 11%, the mean value increased from -0.78 to -0.58 .⁷ Improvements were also registered in maternal education and pathway variables.

<Table 4>

The multivariate analysis results are based on the estimation of the various specifications of Equation 3. As indicated earlier, the dependent variable is a dichotomous variable. Therefore, the models are estimated using logistic regression. The logistic regression model fits the log odds or logits by a linear function of the explanatory variables as follows: $\text{logit}(p_i) = \ln\left[\frac{p_i}{1-p_i}\right] = \alpha + \mathbf{x}'\boldsymbol{\beta}$, where p_i is the probability that the child is stunted or wasted conditional on \mathbf{x} which is a vector of explanatory variables included in Equation 3; $\ln\left[\frac{p_i}{1-p_i}\right]$ is the log odds of the outcome; and α and β are the parameters to be estimated.

4.1 Maternal education and chronic malnutrition

Table 5 reports the log odds of various specifications of chronic malnutrition, stunting. Model 1 is the baseline model with only maternal education included as a primary explanatory variable after controlling for survey year.⁸ Model 2 adds geographic controls (place of residence and regions) to Model 1. Models 3 thru 6 each add a pathway variable to the baseline model after controlling for place of residence, regions and survey year. In this row, Model 3 is the *socioeconomic status* model; Model 4 is the *health - seeking behavior* model; Model 5 is the knowledge model; and Model 6 is the *reproductive behavior* model. Finally, Model 7 presents the full model with all the primary explanatory variables and control variables included.

Maternal education is significant in the baseline model (Model1) where it is controlled only for survey year. Model 2 shows that the addition of geographic controls to the baseline model reduces the education effect while the significance of the education variable remains unchanged. Models 3, 4, 5& 6 show that, except for the health seeking behavior, all other pathways (socioeconomic status, knowledge and reproductive behavior) are significant and the education effect is significant but lower in absolute value when compared to the baseline model. However, in the full model (Model 7), maternal education, socioeconomic status and some reproductive behavior variables are significant. The decline in the significance of some of the pathway variables could be due to multicollinearity either with maternal education or socioeconomic status or both.

The top row of Table 5 shows that the log odds associated with maternal education declining from 0.27 to 0.16 in absolute value. It appears that each level of education decreases the relative probability of stunting by 24 % ($=[1-\exp(\log \text{odds})]*100$) in the baseline model.⁹ The impact declines to 15 % in the full model (see also Table 8). Therefore, the decline of the direct effect from 24% to 15% means that the pathways and geographic controls explained only about 38% of the education effect.

<Table 5>

Referring to the full model (Table 5, Model 7), the important predictors of stunting are, therefore, maternal education, socioeconomic status and reproductive behavior. Socioeconomic status is the most important predictor of stunting as demonstrated by the

magnitude of the coefficient (log odds) and its significance. The likelihood of stunting also increases with child age at a decreasing rate and decreases with maternal age at a decreasing rate. Similar to earlier findings in Africa and other developing countries (e.g. Marcoux, 2002) but in contrast to other studies on Ethiopia (e.g. Girma and Genebo, 2002), the male dummy is significant implying male children are more likely to be stunted than females.

The place of residence dummy (urban =1) is insignificant in all models. The result is expected in multivariate setting due to the fact that the “urban advantage” is captured by other better measures of urban based social and economic amenities (Fotso, 2006). However, some regions (Tigray, Afar, Amhara, Oromyia, Somali & SNNP) are found significantly different from the reference region, Addis Ababa (Model 2). A child from these regions is more likely to be stunted when compared to a child from the reference region, Addis Ababa. The regional variation obtained in Models 2-6 could be due to differences in the level of urbanization.

Finally, the discussion in Section 2 presented significant changes in stunting over time. The results of the full model in Table 5 shows the survey year dummy is insignificant suggesting the absence of difference between 2000 and 2005 when other factors are considered. The change is attributable to changes in other factors including maternal education, socioeconomic status and reproductive behavior.

4.2 Maternal education and acute malnutrition

Table 6 reports logistic regression results of acute malnutrition, wasting. The presentation in Table 6 follows the approach used earlier in Table 5. Therefore, first row in Table 6 demonstrates how the effect of education on wasting changes as new pathway variable is included in the baseline model.

<Table 6>

The comparison of results in Table 5 and Table 4.6 would show the differences and common features of models of chronic and acute malnutrition. First, in Table 6, maternal education is insignificant in all but in the baseline model. Second, both education and pathway variables are also insignificant in the health seeking behavior and knowledge models. However, similar to the chronic malnutrition case, socioeconomic status and

selected reproductive behavior variables are significantly related to acute malnutrition. In addition, geographic and survey year controls are found to have a similar pattern.

4.3. Robustness tests: Alternative sample domains

Table 7 and Table 8 respectively report logistic regression results of the full models of stunting and wasting based on alternative sample domains. The estimations are based on a disaggregated data by survey year and place of residence. Each table incorporates four models. The first two models are for each survey year: 2000 and 2005. The third and the fourth models are for rural and urban children respectively.

<Table 7>

The results in Table 7 are in general similar to the results in Table 5. Accordingly, socioeconomic status is significant in all cases. Education retains its significance in two of the four cases. Health seeking behavior and knowledge are insignificant in most cases. However, health-seeking behavior appears significant in the urban model. Likewise, the results in Table 8 are similar to that of Table 6. In most cases maternal education and its pathways are insignificant.

<Table 8>

4.4 Discussion

Table 9 summarizes the results and compares the effect of maternal education on child nutritional status by model type and measure of malnutrition. The log odds in Table 9 are obtained from the first rows in the previous tables (Tables 5-4). The impact of each level of education on the relative probability of being stunted or wasted is calculated accordingly i.e., $(1-\exp(\log \text{ odds}))$.

<Table 9>

The summary of results in Table 9 indicates that the maximum effect of maternal education on stunting and wasting is observed in the baseline model. It is 24% for stunting and 20% for wasting. In the full model, the effect declines to 15% for stunting and 11% for wasting. The Table also shows that maternal education and its pathways are more relevant to explain stunting than wasting. Except for the baseline model, maternal education is not significant in all other models of wasting. Another important observation is that the direct effect of maternal education is larger in the rural than urban

areas. Each level of maternal education in the rural reduces the relative probability of stunting by 19%. However, it is not significant in the urban areas.

Socioeconomic status is the most important factor of all pathways in mediating the impact of maternal education on child nutritional status. It is significant in all sample categories of stunting and in the national and rural models of wasting.¹⁰ The results imply that policies and programs intended to reduce child malnutrition and hence child mortality would primarily focus on targeting the various key elements of socioeconomic status. Socioeconomic status in this study is measured by the DHS wealth index. Its specific misgivings would make it less amenable to policy. First, the key elements from which the DHS wealth index is constructed are predominantly urban based. Therefore, the index could simply be measuring urbanicity. Second, different elements contribute to the index differently. Therefore, what part of the index is essentially driving the impact on child health requires explanation.

Overall, the results obtained for chronic and acute child malnutrition are in line with earlier related works on other countries including Frost, Forste and Haas (2005) for Bolivia and Webb and Block (2004) for Indonesia. The findings are also robust to changes to sample domains. Disaggregating the sample by survey year and place of residence did not change the results substantially. However, the inability to explain the full effects of maternal education in chronic malnutrition and its erratic relationships with acute malnutrition is an important limitation to the analysis presented in this study. The problem could be due to the presence of other channels that are not considered in this study or measurement error in the variables from which the indexes of the pathways are constructed. Future work on the issue using a different data set and a different country would add more insight in the relationship between maternal education and child nutritional status.

5. Summary and Conclusions

This study models the impact of maternal education and its pathways on chronic and acute child malnutrition in Ethiopia using the 2000 and 2005 Demographic and Health Surveys. The pathways examined in this study are socioeconomic status, maternal health-seeking behavior, maternal knowledge of health and family planning and reproductive

behavior. The logistic models of stunting and wasting are estimated for various sample categories including the national sample, rural sample, urban sample, the 2000 sample and the 2005 sample.

Maternal education works through all pathways except health-seeking behavior. Each level of maternal education reduces the relative probability of being chronically malnourished by 15%. However, no direct effect of maternal education is obtained on acute malnutrition. Overall, maternal education and its pathway explain chronic malnutrition better than acute malnutrition. The claim that maternal education is the single most important predictor of malnutrition would be oversimplification.

Socioeconomic status is the most important pathway linking maternal education and child nutritional status. It is significant in both models of chronic and acute malnutrition. Although girls' education is a high policy priority, it may take time before its direct and indirect impacts substantially improve child health outcomes. Faster results would require direct interventions on key elements of socioeconomic status

¹ According to online databases of United Nations Children's Fund (UNICEF) and United Nations Population Division.

² Although the test of equality of distribution shows that wasting has significantly declined in 2005, the improvement in z-scores is to the right of the cutoff point, -2SD (Figure 2, Panel D2). Therefore, the proportion of wasted children remains the same. However, the results in Table 2 imply that those children who were not wasted in the 2005 sample had a better nutritional status than those in the 2000 sample

³ Estimating structural equation involves endogenous factors such as child health inputs. However, due to difficulties in finding instruments, most studies estimate conditional demand function given by Equation 3 (e.g. Glewwe, 1999; Christiaensen and Alderman, 2004; Webb and Block, 2004 Frost, Forste and Haas, 2005).

⁴ It should be noted that, not all these demographic variables are responsive to maternal education. Therefore, some such as sex and age of child are included as additional control variables.

⁵ Another form of gender biased human capital outcome would result from fertility behavior. Jensen (2002) argues female children may have more siblings than male children as a result of son-preferring differential stopping behavior in fertility, i.e. on average females come from a larger family size where all children are worse off. Therefore, even if there is equal treatment at household level there may be unequal outcome at aggregate level. In some traditional societies son-preferring behavior would result in inequalities in nutritional outcomes (Tarozzi and Mahajan, 2007).


⁶ The pros and cons of the DHS wealth index have been noticed. On the one hand, the DHS surveys are often implemented in countries where income itself may not be the most reliable—or even available—way of measuring socioeconomic status. On the other hand, the index is constructed from urban-based social and economic amenities and may be measuring urbanicity instead of socioeconomic status.

⁷ Percent stunted and HAZ measure long-term malnutrition but in opposite directions. Similarly, percent wasted and WHZ measure short-term malnutrition in opposite directions. The discussion in Section 4.2 is based on z scores. The multivariate analysis is based on the likelihood of being stunted or wasted.


⁸ The pooled (2000 and 2005) data are used in all models. Therefore, all are controlled for a survey year dummy.

⁹ For a unit increase in the j^{th} regressor, the derivation and interpretation is as follows: $\exp(\mathbf{x}'\boldsymbol{\beta})$ increases to $\exp(\mathbf{x}'\boldsymbol{\beta} + \beta_j) = \exp(\mathbf{x}'\boldsymbol{\beta}) * \exp(\beta_j)$. Hence the odds ratio, $[\frac{p_i}{1-p_i}]$ increases by a multiple $\exp(\beta_j)$. Thus, for example, the logit slope parameter of 0.1 means a unit increase in the regressor multiplies the initial odds ratio by $\exp(0.1) = 1.105$. Which is a proportionate increase of 0.105 times the initial odds ratio. Therefore the relative probability of being stunted increases or decreases (depending on the sign of the coefficient in the logit model) by 10.5 percent (Cameron and Trivedi, 2005, p. 470).


¹⁰ The minimum level of statistical significance 5% employed in this study is 5%. However, socioeconomic status is significant at 10% level in the 2000 and 2005 samples. All other pathways are very far from that (Table 7).




(A1) HAZ by Maternal Education: National Sample




(A2) WHZ by Maternal Education: National Sample




(B1) HAZ by Maternal Education: Rural Sample



(B2) WHZ by Maternal Education: Rural Sample




(C1) HAZ by Maternal Education: Urban Sample




(C2) WHZ by Maternal Education: Urban Sample


Figure 1 Density estimates of HAZ and WHZ by Maternal Education




(D1) HAZ by Survey Year: National Sample




(D2) WHZ by Survey Year: National Sample




(E1) HAZ by Survey Year: Rural Sample



(E2) WHZ by Survey Year: Rural Sample



(F1) HAZ by Survey Year: Urban Sample



(F2) WHZ by Survey Year: Urban Sample

Figure 2 Density estimates of stunting and wasting by survey year in 2000 and 2005.

Table 1. Child Malnutrition in Ethiopia by Maternal Education and Place of Residence (in 2000 and 2005)

Background	% of Children suffering from								
	Stunting			Wasting			Underweight		
	2000	2005	% Change	2000	2005	% Change	2000	2005	% Change
Maternal Education									
No Education	52.9	49.1	-7.2	11.4	11.2	-1.8	49.6	41.4	-16.5
Primary	49.1	39.8	-18.9	8.8	10.1	14.8	40.4	32.0	-20.8
Sec. & Higher	32.9	24.0	-27.1	6.7	1.3	-80.6	27.7	13.6	-50.9
Place of Residence									
Urban	42.3	29.8	-29.6	5.5	6.3	14.6	33.7	22.9	-32.1
Rural	52.6	47.9	-8.9	11.1	10.9	-1.8	48.7	39.7	-18.5
Total	51.5	46.5	-9.7	10.5	10.5	0.0	47.2	38.4	-18.6

Source: 2000 and 2005 EDHS (Central Statistical Authority and ORC Macro, 2001&2006).

Note: Stunted =height-for-age z-scores bellow -2; wasted=weight for height z scores bellow -2; and underweight =weight for age z scores bellow -2 (World Health Organization, 1986).

Table 2. Kolmogorov-Smirnov test of equality of distributions of HAZ, WHZ and WAZ by survey year

Test Category	Nutrition Indicator	Sample		
		National	Rural	Urban
Maternal education: Some education Vs No education	HAZ	0.150**	0.077**	0.154**
	WHZ	0.115**	0.067**	0.132**
	WAZ	0.182**	0.093**	0.182**
Place of residence: Urban Vs Rural	HAZ	0.207**	-	-
	WHZ	0.125**	-	-
	WAZ	0.246**	-	-
Survey Year: 2005 Vs 2000	HAZ	0.074**	0.079**	0.090**
	WHZ	0.080**	0.084**	0.090**
	WAZ	0.080**	0.089**	0.123**

Note: The test compares cumulative distributions of each malnutrition indicator in 2000 and 2005;

** =p-value<0.01.

Table 3. Description of variables included in health-seeking behavior and knowledge indexes

Variable	Sample Mean	Factor Loadings	α
Health-seeking behavior Index			
Delivered a baby in modern health facility	0.051	0.491	
Received prenatal from a health professional	0.276	0.546	
Received tetanus injection before birth	0.313	0.490	
Used contraceptive	0.186	0.470	0.718
Knowledge Index			
Has heard of oral rehydration therapy	0.570	0.185	
Knows when in ovulatory cycle can get pregnant	0.408	0.135	
Frequency of reading news paper	0.062	0.377	
Frequency of listening to radio	0.457	0.381	
Frequency of watching TV	0.099	0.439	
Heard of family planning from radio	0.176	0.378	
Heard of family planning from TV	0.026	0.429	
Read about family planning from newspaper	0.019	0.376	0.716

Note: α is a reliability coefficient or Cronbach's alpha which is greater than 0.7 in both cases, which is within acceptable range.

Source: EDHS 2000&2005 (Statistical Authority and ORC Macro, 2001&2006)

Table 4. Descriptive statistics of model variables by survey year

Variables	2000 EDHS				2005 EDHS			
	Min	Max	Mean	SD	Min	Max	Mean	SD
Height for age z-score (HAZ)	-5.99	5.84	-2.06	0.03	-5.98	5.78	-1.77	0.05
Weight for height z-score (WHZ)	-3.99	5.71	-0.78	0.02	-3.99	5.91	-0.58	0.03
Stunted (HAZ<-2SD=1)	0	1	0.51	0.01	0	1	0.46	0.01
Wasted (WHZ<-2SD=1)	0	1	0.11	0.01	0	1	0.11	0.01
Maternal education (categories)	0	5	0.30	0.02	0	5	0.32	0.02
Socioeconomic status (wealth index)	-1.05	3.43	-0.41	0.02	-3.10	3.76	-0.36	0.02
Health seeking behavior	-1.16	4.85	-0.29	0.04	-1.16	3.68	-0.09	0.04
Knowledge of family planning & health	-1.20	9.94	-0.23	0.03	-1.20	9.94	-0.18	0.03
Child is male	0	1	0.51	0.01	0	1	0.51	0.01
Child age (age in months& in logs)	0.00	4.09	3.14	0.01	0.00	4.09	3.10	0.01
Child age squared	0.00	16.76	10.62	0.05	0.00	16.76	10.45	0.07
Mother's age (age in years & in logs)	2.71	3.89	3.36	0.00	2.71	3.89	3.35	0.00
Mother's age squared	7.33	15.14	11.33	0.03	7.33	15.15	11.27	0.03
Birth order >3	0	1	0.51	0.01	0	1	0.53	0.01
Preceding birth interval > 2years	0	1	0.65	0.01	0	1	0.65	0.01

Note: Min=minimum value; Max=maximum value; and SD = standard deviations.

Source: Ethiopian Demographic and Health Survey (EDHS) 2000 and 2005 (Central Statistical Authority and ORC Macro, 2001&2006)

Table 5. Logistic regression results of child stunting (2000 and 2005 pooled)

Variable	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
Maternal education	-0.27** (-7.41)	-0.23** (-5.46)	-0.19** (-4.31)	-0.23** (-4.70)	-0.17** (-3.56)	-0.23** (-5.44)	-0.16** (-2.65)
Socioeconomic status			-0.28** (2.96)				-0.37** (-3.40)
Health-seeking behavior				-0.04 (-1.38)			-0.05 (-1.60)
Knowledge					-0.08** (-2.07)		-0.04 (-0.80)
Reproductive behavior						0.09	0.17**
Sex of child (male=1)						(1.94)	(2.89)
Child age						3.86** (11.18)	3.85** (9.51)
Child age squared						-0.52** (-9.34)	-0.50** (-7.40)
Birth order >3						0.28** (3.58)	0.12 (1.23)
Birth Interval >2years						-0.08 (-1.34)	-0.11 (-1.38)
Mother's age						7.76** (2.63)	6.87* (2.09)
Mother's age squared						-1.21** (-2.78)	-1.07** (-2.21)
Residence (Urban=1)		-0.06 (-0.35)	0.25 (1.15)	-0.01 (-0.05)	0.04 (0.18)	-0.09 (-0.49)	0.34 (1.33)
Survey Year (2005=1)	-0.19** (-3.06)	-0.18** (-2.97)	-0.16** (-2.62)	-0.11 (-1.49)	-0.17** (-2.84)	-0.22 (-3.36)	-0.08 (-1.12)
Constant	0.13 (3.38)	-0.54** (-2.70)	-0.43* (-2.24)	-0.53** (-2.67)	-0.53** (-2.78)	-19.77** (-3.94)	-18.20** (-3.29)
Regional variation? ^a	Yes	Yes	Yes	Yes	Yes	Yes	No ^b
Pseudo R square	0.01	0.02	0.10	0.02	0.02	0.10	0.13
Log Likelihood	-9704	-9632	-9620	-6484	-9556	-8844	-5681
LRChi2	186**	331**	355**	245**	343**	1909**	1764**
N	12463	12463	12463	8572	12350	12463	8498

Note: t-statistics in parentheses; * and ** denote significance at 5% and 1% level respectively; ^a regional variation refers to the 11 geographic regions with Addis Ababa as a reference category; ^b no variation for most regions except for Dire Dawa and Gambella regions where they are found better than the reference category.

Table 6. Logistic regression results of child wasting (2000 and 2005 pooled)

Variable	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
Maternal education	-0.23** (-4.05)	-0.11 (-1.88)	-0.06 (-0.87)	-0.10 (-1.48)	-0.08 (-1.21)	-0.12 (-1.90)	-0.081 (-0.98)
Socioeconomic status			-0.42** (-3.10)				-0.41** (-2.63)
Health-seeking behavior				-0.06 (-1.26)			-0.04 (-0.79)
Knowledge					-0.04 (-0.77)		0.03 (0.58)
Reproductive behavior						0.14 (1.81)	0.20* (2.35)
Sex of child (male=1)							
Child age						3.13** (6.11)	3.44** (5.21)
Child age squared						-0.59** (-6.64)	-0.64** (-5.39)
Birth order >3						0.12 (1.11)	-0.03 (-0.25)
Birth Interval >2years						0.14 (1.55)	0.23* (2.08)
Mother's age						7.84 (1.57)	7.84 (1.54)
Mother's age squared						-1.07 (-1.56)	-1.15 (-1.54)
Residence (Urban=1)		-0.56** (-2.83)	-0.11 (-.45)	-0.45* (-2.07)	-0.52* (-2.43)	-0.54** (-2.63)	-0.01 (-0.04)
Survey Year (2005=1)	-0.01 (-0.16)	-0.07 (-0.79)	-0.04 (-0.78)	-0.08 (-0.82)	-0.07 (-0.76)	-0.06 (-0.67)	-0.05 (-0.51)
Regional variation? ^a	Yes	Yes	Yes	Yes	Yes	Yes	NO
Pseudo R square	0.00	0.01	0.01	0.01	0.01	0.03	0.04
Log Likelihood	-4820	-4790	-4780	-3518	-4756	-4669	-3399
LRChi2	33.3**	94.6**	113.7**	75.4**	94.9**	342**	284**
N	12637	12637	12637	9601	8660	12637	8585

Note: t-statistics in parentheses; * and ** denote significance at 5% and 1% level respectively; ^a regional variation refers to the 11 geographic regions with Addis Ababa as a reference category; ^b no variation for all regions except for Somali Region where the likelihood of child wasting is found larger than the reference category.

Table 7. Logistic regression results of child stunting by survey year and place of residence

	2000	2005	Rural	Urban
Maternal education	-0.14 (-1.76)	-0.18 (-2.04) *	-0.21 (-2.73)**	-0.07 (-0.63)
Socioeconomic status	-0.37 (-2.74)**	-0.48 (-2.33)*	-0.36 (-2.60)**	-0.61 (-2.71)**
Health-seeking behavior	-0.05 (-1.35)	-0.06 (-0.98)	-0.03 (-0.73)	-0.13 (-1.98) *
Knowledge	-0.05 (-0.80)	-0.01 (-0.20)	-0.05 (-0.95)	0.00 (-0.02)
Reproductive behavior				
Sex of child (male=1)	0.16 (2.42) *	0.19 (1.80)	0.23 (3.75)**	-0.38 (-1.92)
Child age	3.70 (7.46) **	4.27 (6.84) **	4.03 (9.94)**	1.91 (1.55)
Child age squared	-0.47 (-5.71) **	-0.58 (-5.42) **	-0.53 (-7.68)**	-0.19 (-0.93)
Birth order >3	0.08 (0.63)	0.26 (1.59)	0.16 (1.66)	-0.08 (-0.19)
Birth Interval >2years	-0.17 (-1.68)	0.02 (0.17)	-0.09 (-1.11)	-0.23 (-1.08)
Mother's age	10.35 (2.61) **	-2.93 (-0.52)	5.89 (1.76)	16.49 (1.27)
Mother's age squared	-1.56 (-2.69) **	0.34 (0.41)	-0.93 (-1.89)	-2.43 (-1.26)
Residence (Urban=1)	0.36 (1.04)	0.26 (1.02)		
Dire Dawa	-0.56 (-2.38) *	-0.25 (-0.65)		-0.56 (-2.64) **
2005=1			-0.07 (-0.84)	-0.37 (-1.67)
Constant	-23.97 (-3.58)**	-1.94 (-0.21)	-16.63 (-2.95)**	-31.07 (-1.48)**
Regional Variation?	No	No	Yes	Yes
Pseudo R square	0.14	0.14	0.14	0.15
Log Likelihood	-4014	-1654	-5079	-550
LRChi2	1254	527	1586	194
N	5971	2527	6600	1450

Note: Note: t-stat in parenthesis; *, ** significant at 5% and 1% level respectively; pooled data is used for rural and urban models; urban regions (Addis Ababa, Dire Dawa and Harari) are excluded in the rural model and the reference category in the rural model is Tigray; the reference category for regional dummies in all other models is Addis Ababa.

Table 8. Logistic regression results of child wasting by survey year and place of residence

	2000	2005	Rural	Urban
Maternal education	0.03 (0.36)	-0.35 (-2.33)*	-0.15 (-1.48)	0.09 (0.51)
Socioeconomic status	-0.34 (-1.82)	-0.63 (-1.88)	-0.41 (-2.05)*	-0.35 (-1.31)
Health-seeking behavior	-0.01 (-0.21)	-0.08 (-0.93)	-0.03 (-0.61)	-0.09 (-0.90)
Knowledge	-0.04 (-0.62)	0.15 (1.54)	0.03 (0.40)	-0.01 (-0.11)
Reproductive behavior				
Sex of child (male=1)	0.18 (1.72)	0.26 (1.71)	0.19 (2.06)*	0.45 (1.31)
Child age	4.46 (6.06)**	1.75 (1.62)	3.40 (4.93)**	4.40 (2.05)*
Child age squared	-0.81 (-6.21)**	-0.34 (-1.74)	-0.63 (-5.10)**	-0.78 (-2.04)*
Birth order >3	0.03 (0.21)	-0.07 (-0.29)	-0.07 (-0.51)	0.26 (0.56)
Birth Interval >2years	0.45 (3.48)**	-0.33 (-1.66)	0.24 (2.08)*	0.21 (0.51)
Mother's age	2.37 (0.39)	21.11 (2.27)*	9.63 (1.82)	-11.16 (-0.52)
Mother's age squared	-0.35 (-0.39)	-3.09 (-2.26)*	-1.40 (-1.81)	1.55 (0.48)
Residence (Urban=1)	-0.12 (-0.36)	0.20 (0.35)		
2005=1			-0.04 (-0.37)	-0.15 (-0.28)
Constant	-12.14 (-1.17)	-40.40 (-2.58)**	-22.97 (-2.57)**	11.37 (0.33)
Regional variation?	No	No	Yes	Yes
Pseudo R square	0.05	0.06	0.04	0.10
Log Likelihood	-2405	-947	-3149	-224
LRChi2	250	115	223	50
N	6058	2527	6682	1454

Note: Note: t-stat in parenthesis; *, ** significant at 5% and 1% level respectively; pooled data is used for rural and urban models; urban regions (Addis Ababa, Dire Dawa and Harari) are excluded in the rural model and the reference category in the rural model is Tigray; the reference category for regional dummies in all other models is Addis Ababa.

Table 9. The effect of maternal education and pathway variables on stunting and wasting by model type

Model Type	Chronic Malnutrition (Stunting)				Acute Malnutrition (Wasting)			
	log odds	1-exp (log odds)	Pathway		log odds	1-exp (log odds)	Pathway	
			Education variable significant ?	Education variable significant ?			Education variable significant ?	Education variable significant ?
Baseline	-0.27	0.24	Yes	–	-0.23	0.20	Yes	–
Geographic	-0.23	0.21	Yes	Yes	-0.11	0.10	No	Yes
SES	-0.19	0.17	Yes	Yes	-0.06	0.05	No	Yes
Health-seeking	-0.23	0.21	Yes	No	-0.10	0.10	No	No
Knowledge	-0.17	0.16	Yes	Yes	-0.08	0.08	No	No
Reproductive	-0.23	0.21	Yes	Yes	-0.12	0.11	No	Yes
Full Model								
National	-0.16	0.15	Yes	Yes ^a	-0.12	0.11	No	Yes ^a
Rural Sample	-0.21	0.19	Yes	Yes ^a	-0.15	0.14	No	Yes ^a
Urban Sample	-0.07	0.07	No	Yes	-0.09	0.09	No	No
2000 Sample	-0.14	0.13	No	Yes	0.03	-0.03	No	No
2005 Sample	-0.18	0.16	Yes	Yes	-0.35	0.30	Yes	No

Note: SES is socioeconomic status; ^a some pathways only (mostly SES).

References

- Alderman, Harold , Jere R. Behrman, Victor Lavy and D. Menon, (2001), "Child health and school enrollment: A longitudinal analysis," *The Journal of Human Resources*, 36(1): 185-205.
- Behrman, Jere R., (1988), "Intrahousehold allocation of nutrients in rural India: Are boys favored? Do parents exhibit inequality aversion?" *Oxford Economic Papers*, 40(1): 32-54.
- Behrman, Jere R. (1997), "Intrahousehold distribution and the family," in Mark Rosenzweig and Oded Stark (eds), *Handbook of Population and Family Economics*, Chapter 4. Vol. 1A., Elsevier.
- Behrman, Jere R., and Anil B. Dolalihar (1990), "The intrahousehold demand for nutrients in rural South India: individual estimates, fixed effects and permanent income."
- Black, Robert E., Saul .S. Morris, and Jennifer Bryce, (2003), "Where and why are 10 million children dying every year?" *Lancet*, 361(9376):2226-2234.
- Caldwell, J.C., (1979), "Education as a factor in mortality decline: An examination of the Nigerian data," *Population Studies*, 33(3): 395-413.
- Cameron, Colin A., and Pravin K. Trivedi, (2005), *Microeconometrics: Methods and Applications*, Cambridge University Press.
- Central Statistical Authority (CSA), Ethiopia and ORC Macro, (2001), Ethiopia Demographic and Health Survey 2000. Addis Ababa, Ethiopia and Calverton, Maryland USA: CSA and ORC Macro.
- Central Statistical Authority (CSA), Ethiopia and ORC Macro, (2006), Ethiopia Demographic and Health Survey 2005, Addis Ababa, Ethiopia and Calverton, Maryland USA: CSA and ORC Macro.
- Christiaensen, Luc, and Harold Alderman, (2004), "Child malnutrition in Ethiopia: Can maternal knowledge augment the role of income," *Economic Development and Cultural Change*, 52(2):287-312.
- Desai, Solande, and Soumya Alva, (1998), "Maternal education and child health: Is there a strong causal relationship?" *Demography*, 35(1): 71-81.
- Fotso, Jean-Christophe, (2006), "Child health inequities in developing countries: differences across urban and rural areas," *International Journal for Equity in Health*, 5(9).
- Fongillo, Edward Jr., Mercedes de Onis, and Kathleen M.P. Hanson, (1997), "Socioeconomic and demographic factors are associated with worldwide patterns of stunting and wasting," *Journal of Development* 127(12): 2302-2309.
- Girma, Woldesenbet, and Timotiws Genebo, (2002), *Determinants of Nutritional Status of Women and Children in Ethiopia*. Calverton, Maryland, USA: ORC Macro.
- Glewwe, Paul, (1999), "Why does mother's schooling raise child health in developing countries? Evidence from Morocco." *Journal of Human Resources*, 34(1). 124-159.

- Jensen, R., (2002), "Equal Treatment, Unequal Outcomes? Generating Gender Inequality Through Fertility Behavior," Working Paper, JFK School of Government, Harvard.
- Jeyaseelan, L., and M. Lakshman, (1997), "Risk factors for malnutrition in South Indian children," *Journal of Biosocial Science*, 29(1): 93-100.
- Marcoux, Alain, (2002), "Sex differentials in Undernutrition: A look at survey evidence," *Population and Development Review*, 28(2): 275-284.
- Opinion Research Company (ORC) Macro, (2001), *Nutrition of young children and mothers in Ethiopia 2000*. Africa Nutrition Chartbooks. Calverton, Maryland, USA: ORC Macro.
- Pitt, M. Mark. and Mark R. Rosenzweig, (1985), "Health and nutrient Consumption Across and Within Farm Households," *The Review of Economic Statistics*, 67(2): 212-223.
- Pongou, Roland, Majid Ezzati and Joshua Salomon, (2006) "Household and community socioeconomic and environmental determinants of child nutritional status in Cameroon," *BMC Public Health*, 6: 98. <http://www.biomedcentral.com/1471-2458/6/98>.
- Smith, Lisa, and Lawrence Haddad, (2000), Experiencing Child Malnutrition in Developing Countries: A Cross-Country Analysis. Research Report no. 111, International Food Policy Research Institute, Washington D.C.
- Sommerfelt, A. Elizabeth, and S. Kathryn, (1994), *Children's nutritional status*. DHS Comparative Studies No. 12. Calverton, Maryland, USA: ORC Macro.
- Strauss, John, and Duncan Thomas, (1998), "Health, nutrition and economic development," *Journal of Economic Literature*, 36(2): 766-817.
- Tarozzi, Alessandro and Aprajit Mahajan, (2007), "Child nutrition in India in the nineties" *Journal of Economic Development and Cultural Change*, 55(3): 441-486.
- Thomas, Duncan, Victor Lavy and John Strauss, (1996), "Public Policy and Anthropometric Outcomes in the Côte D'Ivoire," *Journal of Public Economics*, 61(2): 155-192.
- Variyam, Jayachandran N., James Blaylock, Biing-Hwan Lin, Katherine Ralston, and David Smallwood, (1999), "Mother's nutrition knowledge and children's dietary intakes," *American Journal of Agricultural Economics*, 81(2): 373-384.
- Webb, Patrick and Steven Block, (2004), "Nutrition information and formal schooling as inputs to child nutrition," *Economic Development and Cultural Change*, 55(2).
- Wolfe, Barbara, and Jere Behrman, (1983), "Is income over-rated in determining adequate nutrition?" *Economic Development and Cultural Change*, 31(4): 525-549.
- World Health Organization(WHO) Working Group, (1986), "Use and interpretation of anthropometric indicators on nutritional status," *Bulletin of the World Health Organization*, 64(6): 929-941.