Western Michigan University ScholarWorks at WMU

Dissertations

Graduate College

12-2015

Rank Based Procedures for Ordered Alternative Models

Yuanyuan Shao Western Michigan University, shaoyy_mei@yahoo.com

Follow this and additional works at: https://scholarworks.wmich.edu/dissertations

Part of the Statistics and Probability Commons

Recommended Citation

Shao, Yuanyuan, "Rank Based Procedures for Ordered Alternative Models" (2015). *Dissertations*. 1197. https://scholarworks.wmich.edu/dissertations/1197

This Dissertation-Open Access is brought to you for free and open access by the Graduate College at ScholarWorks at WMU. It has been accepted for inclusion in Dissertations by an authorized administrator of ScholarWorks at WMU. For more information, please contact wmu-scholarworks@wmich.edu.





RANK BASED PROCEDURES FOR ORDERED ALTERNATIVE MODELS

by Yuanyuan Shao

A dissertation submitted to the Graduate College in partial fulfillment of the requirements for the degree of Doctor of Philosophy Statistics Western Michigan University December 2015

Doctoral Committee:

Joseph W. McKean, Ph.D., Chair Jeffery T. Terpstra, Ph.D. Jung Chao Wang, Ph.D. Bradley E. Huitema, Ph.D.

RANK BASED PROCEDURES FOR ORDERED ALTERNATIVE MODELS

Yuanyuan Shao, Ph.D.

Western Michigan University, 2015

The ordered alternatives in a one-way layout with k ordered treatment levels are appropriate for many applications, especially in psychology and medicine. There is extensive literature in this area, and many parametric and nonparametric approaches have been introduced. Abelson-Tukey (AT) test is a frequently used parametric method. Its coefficients provide an ideal way of combining means for the purpose of detecting a monotonic relationship between the independent and dependent variables. The AT method, though, is not robust. Furthermore, our initial empirical studies show that it is not more powerful than the Jonckheere-Terpstra (JT) and the Hettmansperger-Norton (HN) nonparametric tests at normal errors for moderate sample sizes. Theses nonparametric tests, unlike the AT test, are not easily extended to general linear and mixed models.

We have developed a rank-based procedure which has the same optimal efficacy properties as the HN procedure for the ordered and umbrella alternative problems, including the unknown peak problem. It is a rank-based procedure and is easily extended to linear, mixed and covariance models. The procedure can utilize general score functions. ©2015 Yuanyuan Shao

ACKNOWLEDGMENTS

I would like to give my deepest thanks to my advisor Dr. Joseph W. McKean for his guidance and patience. I would also express my gratitude to the committee members, Dr. Jung-Chao Wang, Dr. Jeff Terpstra, and Dr. Bradley E. Huitema.

I would also like to thank my husband, Qiang. This dissertation could not be done without his encourgement and support.

Yuanyuan Shao

TABLE OF CONTENTS

A	CKN	OWLI	EDGMENTS	ii
LI	ST C	OF TA	BLES	vi
LI	ST C	OF FIC	GURES	viii
C	HAP	TER		1
1	INT	RODU	UCTION	1
	1.1	Overv	iew of k -sample Location Problem $\ldots \ldots \ldots \ldots \ldots \ldots \ldots$	2
		1.1.1	Basic One-way Notation	2
		1.1.2	Hypotheses	3
	1.2	Detail	s on Some Existing Methods	4
		1.2.1	Distribution-free Nonparametric Tests	4
		1.2.2	Parametric-based Test	8
	1.3	Exam	ples	10
2	мо	TIVA	ΓΙΟΝ	12
	2.1	Initial	Study Results	12
		2.1.1	Empirical α Levels	13
		2.1.2	Empirical Powers	13
	2.2	Robus	t Abelson-Tukey (RAT) Test	20
		2.2.1	Rank-based Estimator	20
		2.2.2	General Rank Scores	21
		2.2.3	The RAT Test	24
	2.3	Simula	ation Results	29

3 NEW METHODS

3	NE	W METHODS	33
	3.1	Robust Abelson-Tukey with HN Weights (RAThn) Test	34
		3.1.1 The RAThn Test Statistic	35
		3.1.2 Consistency and Asymptotic Power	36
	3.2	Robust Shao-McKean (RSM) Test	37
		3.2.1 The RSM Test Statistic	38
		3.2.2 Consistency and Asymptotic Power	46
	3.3	The RSM with Least Square Estimates (RSM_{LS}) Test $\ldots \ldots \ldots$	48
		3.3.1 The RSM_{LS} Test Statistic	48
		3.3.2 Consistency and Asymptotic Power	50
	3.4	Comparisons of New Methods	51
	3.5	Simulation Results	52
		3.5.1 With $N(0, 1)$ Underlying Distribution	53
		3.5.2 With $CN(0.25, 100)$ Underlying Distribution	54
		3.5.3 With $SCN(0.25, 1, 100)$ Underlying Distribution	55
		3.5.4 With $GF(1,.1)$ Underlying Distribution	57
	3.6	Example	60
4	MI	XED MODELS WITH RANDOM BLOCK EFFECT	62
	4.1	Simple Mixed Models	63
	4.2	The Page Test	64
	4.3	New Methods	65
		4.3.1 Abelson Tukey Test with Randomized Blocks(ATb)	66
		4.3.2 Joint Rank Estimator	67
		4.3.3 Nonparametric Methods	67
	4.4	Simulation Studies	70
5	UN	KNOWN PEAK	77
	5.1	The Hettmansperger and Norton Method	77
	5.2	New Methods	78
		5.2.1 RAThn	78

		5.2.2 RSM	78
		5.2.3 RSMls	79
	5.3	Simulation Studies	79
6	ON	E-WAY LAYOUT WITH COVARIATES	86
	6.1	The Covariance Model	87
	6.2	The AT Test	87
	6.3	The RAThn Test	88
	6.4	Simulation Studies	89
		6.4.1 With Normal Underlying Distribution	90
		6.4.2 With $CN(0.25, 100)$ Underlying Distribution	90
		6.4.3 With $SCN(0.25, 1, 100)$ Underlying Distribution	92
7	CO	NCLUSIONS AND FUTURE WORK	94
	7.1	Conclusions	94
	7.2	Future Work	97

LIST OF TABLES

1	Example: Number of parts produced	10
2	Test results	10
3	Initial simulation on empirical α s	13
4	Initial simulation on empirical power under the $Normal$ distribution	15
5	Initial simulation on the empirical power under SCN distribution	19
6	Empirical α s at nominal levels	29
7	Empirical power at nominal levels	29
8	Empirical power levels under the Normal distribution	30
9	Empirical power levels under the SCN distribution	31
10	Empirical power under a <i>Normal</i> distribution for all methods	53
11	Empirical power under $CN(0.25, 100)$ with Wilcoxon scores	54
12	Empirical power under $CN(0.25, 100)$ with bent4 scores	55
13	Empirical power under $SCN(0.25, 1, 100)$ with Wilcoxon scores	56
14	Empirical power under $SCN(0.25, 1, 100)$ with bent1 scores	57
15	Empirical power under $GF(1, 0.1)$ with Wilcoxon scores	58
16	Empirical power under $GF(1, 0.1)$ with logGF scores	59
17	Empirical power under $GF(1, 0.1)$ with bent1 scores	59
18	Test results-new methods	61
19	Empirical α s and power(10x3)	73
20	Empirical α s and power (10x4)	74
21	Empirical α s and power (10x7)	75
22	Empirical α s and power (15x7)	76
23	Unknown peaks estimation (clear peak $N(\mu_i, 1)$)	81
24	Unknown peaks estimation (flat peak under $N(\mu_i, 1)$)	82

25	Unknown peaks estimation (clear peak $CN(0.25, \mu_i, 100)$.)	83
26	Unknown peaks estimation (flat peak $CN(0.25, \mu_i, 100))$	84
27	Empirical α and power under <i>Normal</i> distribution	90
28	Empirical α and power under $CN(0.25,100)$ distribution with Wilcoxon	
	scores	91
29	Empirical α and power under $CN(0.25,100)$ distribution with bent4 scores.	92
30	Empirical α and power under SCN distribution with Wilcoxon scores	92
31	Empirical α and power under SCN distribution with bent1 scores	93

LIST OF FIGURES

1	Boxplot of example data	11
2	Plot of the initial study on power curves under a $Normal$ distribution	15
3	Plot of the initial study on power curves under $t_{(10)}$	16
4	Plot of the initial study on power curves under $t_{(8)}$	16
5	Plot of the initial study on power curves under $t_{(5)}$	17
6	Plot of the initial study on power curves under $t_{(3)}$	17
7	Plot of the initial study on power curves under $t_{(2)}$	18
8	Plot of the initial study on power curves under $t_{(1)}$	18
9	Plot of the initial study on power curves under SCN distribution	19
10	Plots of score functions.	23
11	Plot of power curves under a <i>Normal</i> distribution	30
12	Plot of power curves under the SCN distribution	31
13	Plot of power curves under a $Normal$ distribution for all methods	53
14	Plot of power curves under CN distribution with Wilcoxon scores	54
15	Plot of power curves under CN distribution with bent4 scores	55
16	Plot of power curves under SCN distribution with Wilcoxon scores. $\ .$.	56
17	Plot of power curves under SCN distribution with bent1 scores	57
18	Plot of power curves under GF distribution with Wilcoxon scores	58
19	Plot of power curves under GF distribution with logGF scores	59
20	Plot of power curves under GF distribution with bent1 scores	60
21	Plots of clear peak data with different peaks.	80
22	Plots of flat peak data with different peaks.	80
23	Plot of power curves under Normal distribution	90
24	Plot of power curves under CN distribution (Wilcoxon scores)	91

25	Plot of power curves under CN distribution (bent4 scores)	91
26	Plot of power curves under SCN distribution (Wilcoxon scores)	92

27 Plot of power curves under SCN distribution (bent1 scores). 93

CHAPTER 1

INTRODUCTION

This research is on nonparametric methods of testing for k-sample location problem with ordered alternatives. The null hypothesis of interest is that there are no differences in locations (or no treatment effects), under which all k samples can be treated as a sample from one population. The alternatives considered here correspond to a trend in the locations. For example, consider a randomized group experiment with five levels of the independent variable, where the treatments are doses of a drug (say, 10, 20, 30, 40 and 50 mg). With increasing dose level, or concentration the performance of the treatment tends to improve. This is the alternative of interest.

In some cases, the researchers form the research hypothesis as either an increasing or decreasing trend based on the preliminary information they have obtained, and such a hypothesis is known as the ordered alternative hypothesis. To test such a hypothesis, several test procedures are available. When the alternative is that at least two of the k underlying distributions have different centers, the well-known test procedure is the Kruskal and Wallis (Kruskal and Wallis, 1952). When the alternative is that at least one strict inequality follows for all centers, Terpstra (1952) and Jonckheere (1954) were the first, and independently suggesting the same test. Abelson and Tukey (Abelson and Tukey, 1963), Puri (Puri, 1965), Odeh (Odeh, 1971), Archambault et al. (Archambault et al., 1977), Hettmansperger and Norton (Hettmansperger and Norton, 1987), Büning (Büning, 1999), and McKean et al.(McKean et al., 2001) have also developed methods for these hypotheses. When the treatment effect has a monotonic increasing trend and changes in direction after reaching a peak, it is called an umbrella trend test. Mack and Wolfe (Mack and Wolfe, 1981), Hettmansperger and Norton (Hettmansperger and Norton, 1987), Shi (Shi, 1988), Chen and Wolfe (Chen and Wolfe, 1990), Chen (Chen, 1991), Pan (Pan, 1996), Terpstra and Magel (Terpstra and Magel, 2003), Kössler (Kössler, 2006) and Alvo (Alvo, 2008) discuss nonparametric tests for umbrella alternatives.

In chapter 2, Monte Carlo simulation results of an initial study of the α levels and powers of some of these existing methods are discussed as motivation for this study. In chapter 3, two new distribution-free tests and a parametric-bsed test are proposed, and their properties are developed. A simulation study involving these new procedures, comparing them with those mentioned in chapter 2, is then discussed. In chapter 4, new methods of unknown peak problems are proposed. In chapter 5, new methods for mixed models with random block effects are proposed and simulation results are discussed.

1.1 Overview of k-sample Location Problem

1.1.1 Basic One-way Notation

Let $Y_{1j}, Y_{2j}, \ldots, Y_{n_jj}, j = 1, \ldots, k$ represent a random sample of the response of the *jth* group with sample size n_j having common cumulative distribution function (cdf) $F_j(y)$, where $F_j(y) = F(y - \theta_j)$, where F is an absolutely continuous distribution function. Assume that θ_j $(j = 1, \ldots, k)$ is a location parameter for the population. In the following, we assume that F is twice continuously differentiable on $(-\infty, \infty)$. There are $n = \sum_{j=1}^k n_j$ observations in all. All responses are assumed to be independent of one another. Hence our full model design is the same as the one-way analysis of variance design, that is, the response Y_{ij} follows the linear model,

$$Y_{ij} = \theta_j + \epsilon_{ij}, \ j = 1, \dots, \ k, \ i = 1, \dots, \ n_j,$$
 (1.1)

 Y_{ij} is the *ith* observation on response variable for the *jth* treatment level; θ_j is the *jth* location parameter; ϵ_{ij} are independent and identically distributed (iid) with continuous distribution function F(x) and probability density function f(x). The null hypothesis

of interest is that there is no differences in locations among k populations, or there is no difference in treatment effects, and we write it as

$$H_0: \theta_1 = \theta_2 = \dots = \theta_k. \tag{1.2}$$

Under the null hypothesis all k samples can be treated as one sample from a single population.

1.1.2 Hypotheses

In this thesis we are interested in several alternatives which are given by the following.

Ordered Alternatives

According to the natural labeling of the treatments, the appropriate alternatives could be set as increasing (or decreasing) treatment effects. The monotone alternative hypotheses are of the form:

$$H_1: \theta_1 \le \theta_2 \le \dots \le \theta_k, \text{ (with at least one strict inequality)}, \tag{1.3}$$

or,

$$H_2: \theta_1 \ge \theta_2 \ge \cdots \ge \theta_k$$
, (with at least one strict inequality). (1.4)

Both sets of these alternatives are called ordered alternatives.

Umbrella Alternatives

In some experiments, the performance of the treatment might tend to increase with increasing levels, but after some point higher levels tend to diminish the performance. This is called an umbrella pattern and can be written as:

$$H_3: \theta_1 \le \theta_2 \le \dots \le \theta_p \ge \theta_{p+1} \ge \dots \ge \theta_k. \tag{1.5}$$

This is an umbrella alternative with peak p, where $p \in \{1, 2, ..., k\}$. Thus the ordered alternatives 1.3 are special umbrella alternatives with peak at k.

1.2 Details on Some Existing Methods

Several methods have been proposed for ordered alternatives, including the Terpstra-Jonckheere test which was proposed by Terpstra (1952) and Jonckheere (1954); the Abelson-Tukey test which was proposed by Abelson and Tukey (1963); the Hettmansperger and Norton procedure (1987); and the bootstrap based Spearman approach which was proposed by McKean, Naranjo and Huitema (2001).

These tests are classified into nonparametric (distribution free) and parametric tests.

We next describe these tests.

1.2.1 Distribution-free Nonparametric Tests

Jonckheere-Terpstra(JT) Test

Jonckheere (Jonckheere, 1954) and Terpstra (Terpstra, 1952) were the first to consider ordered alternative rank tests. They independently proposed the following procedure.

The Jonckheere-Terpstra statistic, J is

$$J = \sum_{u=1}^{v-1} \sum_{v=2}^{k} U_{uv},$$

where

$$U_{uv} = \sum_{i=1}^{n_u} \sum_{j=1}^{n_v} \phi(Y_{iu}, Y_{jv}), 1 \le u < v \le k,$$

and $\phi(a, b) = 1$ if a < b, 0 otherwise. Note that U_{uv} is the Mann-Whitney statistic (Mann and Whitney, 1947) for testing the difference between populations u and v. It can be shown that the JT statistics can be described asymptotically as a linear combination of Chernoff and Savage (Chernoff and Savage, 1958) statistics.

For a specified level of significance α , the decision rule is,

Reject H_0 (1.2) in favor of H_A (1.3) if $J \ge j_{\alpha}$; otherwise do not reject,

where the constant j_{α} is chosen to make the type I error probability equal to α .

Under H_0 , J is distribution-free. Although some tables for the null distribution exist, usually the asymptotic test is used. Under H_0 the expected value and variance of J are

$$E_0(J) = \frac{n^2 - \sum_{j=1}^k n_j^2}{4}$$

$$\operatorname{var}_{0}(J) = \frac{n^{2}(2n+3) - \sum_{j=1}^{k} n_{j}^{2}(2n_{j}+3)}{72}.$$

The standardized version of J is

$$J^* = \frac{J - \mathcal{E}_0(J)}{\sqrt{\operatorname{var}_0(J)}} = \frac{J - \left[\frac{n^2 - \sum_{j=1}^k n_j^2}{4}\right]}{\left[\frac{n^2(2n+3) - \sum_{j=1}^k n_j^2(2n_j+3)}{72}\right]^{1/2}}$$

When H_0 is true and as $\min(n_1, n_2, \ldots, n_k)$ tends to infinity, J^* has an asymptotic N(0, 1) distribution (Hollander and Wolfe, 1999).

Thus, the asymptotic test is,

Reject
$$H_0$$
 (1.2) in favor of H_A (1.3)
if $J^* \ge z_{\alpha}$; otherwise do not reject

The Spearman (SP) Test

The Spearman correlation coefficient (Spearman, 1904) (r_s) is well known in psychology. For the ordered alternative problems, the Spearman rho (ρ_s) statistic is used to test the correlation between the ordered levels of the treatment and the dependent variable $(Y_{ij}s)$. If the null hypothesis (1.2) is true, then the population Spearman correlation $\rho_s = 0$; if the ordered alternative hypothesis (1.3 or 1.4) is true, then $\rho_s \neq 0$ (McKean et al., 2001). The Spearman rank correlation coefficient is defined by

$$r_s = \frac{12\sum_{i=1}^n \left[(R_i - \frac{n+1}{2})(S_i - \frac{n+1}{2}) \right]}{n(n^2 - 1)}$$
$$= 1 - \frac{6\sum_{i=1}^n D_i^2}{n(n^2 - 1)},$$

where R_i denotes the rank of X_i among all X; S_i denotes the rank of Y_i among all Y; and $D_i = S_i - R_i$, for i = 1, ..., n.

Under H_0 , the expected value and variance of r_s are

$$\mathcal{E}_0(r_s) = 0,$$

and

$$\operatorname{var}_0(r_s) = \frac{1}{n-1},$$

respectively. The standardized version of r_s is

$$r_s^* = \frac{r_s}{\{\operatorname{var}_0(r_s)\}^{1/2}} = (n-1)^{1/2} r_s.$$

When H_0 is true, r_s^* has an asymptotic N(0, 1) distribution (Hollander and Wolfe, 1999).

The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3) if $r_s^* \ge z_{\alpha}$; otherwise do not reject.

The Bootstrap Spearman Test (BSP)

This procedure uses bootstrap methodology to independently sample n observations with replacement from the observed data Y, and randomly assign these observations in k groups. The Spearman correlation coefficient is then computed for this bootstrap sample. After sampling B (# of bootstraps, which should be large, 10,000 or more) times, the proportion of the sampled Spearman correlation coefficient that are equal to or greater than the observed Spearman correlation coefficient is the observed one-tailed p value (McKean et al., 2001). The statistic used in the bootstrap Spearman

test (BPS) is

$$\bar{R}_n = \sum_{i=1}^k i\bar{R}_{in}.$$

This is Spearman's rho statistic (Tyron and Hettmansperger, 1973; Kendall and Stuart, 1961), where \bar{R}_{in} is the average of the ranks of the items in the *i*th sample. And \bar{R}_n can be written as

$$\bar{R}_n = \frac{1}{n} \sum_{u=1}^{v-1} \sum_{v=2}^k (v-u) U_{uv} + \frac{k(n+1)(k+1)}{4}.$$

The BPS procedure provides:

- 1. A test for ordered alternatives to the null hypothesis of equal population medians,
- 2. A point estimate of the magnitude of monotonic association between the ordered treatment levels and the dependent variable,
- 3. A confidence interval for the measure of the magnitude of association.

Note that the confidence interval is obtained from a second bootstrap which is resampled within each of the original samples.

Hettmansperger and Norton (HN) Test

The Hettmansperger and Norton (HN) test is based on a linear combination of two-sample Chernoff-Savage type statistics computed from k(k-1)/2 pairs of samples. The optimal weights are obtained so as to maximize the Pitman efficacy such that this procedure regarded the tests based on linear contrasts and derived the most efficient rank test for a given pattern, as discussed below (Hettmansperger and Norton, 1987). The test statistic is

$$V^* = \left(\frac{12}{n+1}\right)^{1/2} \frac{V}{\left(\sum a_j^2 / \lambda_j\right)^{1/2}}$$

where $V = \sum_{j} a_j \bar{R}_j$, $\bar{R}_j = \frac{1}{n_j} \sum_{j} R_{ij}$, R_{ij} is the rank of Y_{ij} in the combined samples, $\lambda_j = \frac{n_j}{n}$.

The Pitman efficacy of the test is maximized by $a_j = \lambda_j (c_j - \bar{c}_w)$, where c_j is a set of constants, which specifies the pattern of the alternative, $\bar{c}_w = \sum \lambda_j c_j$, and

 $\sum_{j} a_{j} = 0$. Often $c_{j} = j$ is used, which is optimal under the alternative of an increasing trend and equally spaced means.

If $H_0: \theta_1 = \cdots = \theta_k$ is true,

$$\mathcal{E}_0(V) = 0$$

$$\operatorname{Var}_{0}(V) = [(n+1)/12] \sum \lambda_{i} (c_{j} - \bar{c_{w}})^{2},$$

(Hettmansperger and Norton, 1987). The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3) if $V^* \ge z_{\alpha}$; otherwise do not reject.

They also provided statistics that are based on pairwise ranking and are equivalent in a Pitman efficacy sense to V^* . The statistic is

$$U^* = \sum \sum_{i < j} (d_j - d_i) W_{ij},$$

where $d_j = \lambda_j (c_j - \bar{c}_w)$, $W_{ij} = n_i^{-1} n_j^{-1} U_{ij}$. Under the null hypothesis $H_0: \theta_1 = \cdots = \theta_k$, $E_0(U^*) = 1/2 \sum_{i < j} (d_j - d_i)$ and

$$\operatorname{var}_{0}(U^{*}) = \frac{1}{12} \left\{ \sum_{i < j} \frac{(d_{j} - d_{i})^{2}}{n_{i}n_{j}} + k \sum_{i < j} \left(\frac{d_{j}}{n_{j}} - \frac{d_{i}}{n_{i}}\right) (d_{j} - d_{i}) \right\}.$$

The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3) if $(U^* - \mathcal{E}_0(U^*)) \operatorname{var}^{-1/2}(U^*) \ge z_{\alpha}$; otherwise do not reject.

1.2.2 Parametric-based Test

Abelson-Tukey (AT) Test

This is a frequently used parametric method. The contrast coefficients provide an ideal way of combing means for the purpose of detecting a monotonic relationship between the independent and dependent variables (Abelson and Tukey, 1963). The h_j were selected in an attempt to maximize the minimum correlation between h_j and the unknown X_j (dose for the *jth* group), considering all possibilities under the monotonicity assumption.

The test statistic is defined as

$$t = \frac{\sum_{j=1}^{k} h_j \bar{Y}_j}{\sqrt{MSE\left[\sum_{j=1}^{k} \frac{h_j^2}{n_j}\right]}},$$

where

- 1. h_j is the Abelson-Tukey maximum contrast coefficient associated with treatment j, $h_j = \sqrt{(j-1)[1-(j-1)/k]} - \sqrt{j(1-j/k)};$
- 2. \bar{Y}_j is the *j*th sample mean;
- 3. *MSE* is the within-group mean square from the ANOVA (one-way least squares (LS) fit) on Y;
- 4. n_j is the sample size associated with the *j*th treatment; and
- 5. t is the test statistic.

The AT test selects the h_j s so that the minimum value of the correlation coefficient is compatible with the restrictions on X_j s are maximized, where $\sum h_j = 0$, and provides a standardized effect size measure for the maxmin contrast. If we write $\boldsymbol{\theta} = \boldsymbol{\alpha} + \boldsymbol{X}\boldsymbol{\beta}$, then the problem is the same as to detect $\boldsymbol{\beta}$. Thus, we can use the linear model to test any contrasts of the $\boldsymbol{\theta}$ s. It's easy to extend the test to a single covariate, or multiple covariates and mixed effects cases, and the test will stay same.

However, the AT test is based on a least squares fit, thus, its result is not robust when the underlying distribution of the errors is far from normal distribution (either skewed or heavy tailed). This motivates us to propose a new method, using a rank-based fit and having all the advantages of the AT test.

1.3 Examples

Hundal (1969) described a study on the effects of knowledge of performance on workers whose job was a repetitive task. It was thought that with increased degree of knowledge of performance, the output would be increased. Eighteen male workers were selected and randomly divided into three groups: A (control group), B (treatment group) and C (treatment group). Workers in group A received no information about their performance. Workers in group B received some information about their performance. Workers in group C received detailed information about their performance. The responses indicated were the number of parts each worker processed in the experimental period. Note that to show the robustness of nonparametric tests, the third value in group B has been changed from 54 to 74.

group A	group B	group C
40	38	48
35	40	40
38	74	45
43	44	43
44	40	46
41	42	44

Table 1: Example: Number of parts produced

Let μ_i denote the mean output of group *i*, then the alternative hypothesis is $H_A: \mu_A \leq \mu_B \leq \mu_C$ with at least one strict inequality. Figure 1 shows the boxplot of three groups of data. There is an outlier in group B. The test statistics and *p* values of all mentioned methods are provided in Table 2. The AT test result is affected by the outlier. All methods reject the null hypothesis except the AT test with a *p*-value 0.1989.

Table 2: Test results					
methond	test statistic	p-value			
JT	78	0.0251			
SP	2.2046	0.0212			
BSP	0.4827	0.0220			
HN	1.9737	0.0242			
AT	0.8703	0.1989			



Figure 1: Boxplot of example data

CHAPTER 2

MOTIVATION

A Monte Carlo simulation method was conducted to compare the performance of the JT, SP, HN and AT procedures. The intent of the study is to compare these methods over a wide range of tail weights for the random error distributions. The normal distribution and Student t-distribution with degrees of freedom 10, 8, 5, 3, 2, and 1 (Cauchy) were selected. This is a range from light to very heavy tailed distributions. A skewed contaminated normal distribution was also used which is a mixing of a standard normal distribution with a normal distribution with mean 1 and variance 100, and with the contamination rate at 0.25. The procedures are location and scale equivariant, so standard (convenient) forms of the distributions are used. For this initial investigation we set k = 5 and used group sample sizes of 5, so there were 25 subjects.

A situation consists of a design, a hypothesis and an error distribution. For each situation we ran 10,000 simulations.

Initial study results are shown in section one. A new method, Robust Abelson-Tukey (RAT) test, is proposed in section two. All the simulation results are discussed in section three.

2.1 Initial Study Results

In the first part, the empirical α levels are obtained by repeating this process 10,000 times under the null hypothesis, and computing the proportions of all the *p*-values, which are less than or equal to the chosen alpha level (.05 and .1) on each

procedure.

In the second part, the empirical power values are obtained by repeating the process, similar as in section one, but under the specified alternative hypothesis, 10,000 times. Suppose the center of each group is equally spaced, thus the only difference is an estimate of spacing between the centers being added to the data set of each group.

2.1.1 Empirical α Levels

Table 3 shows the simulation results of the empirical α s at norminal levels 10% and 5% under different population distributions.

Since SP, JT and HN are distribution free, this null simulation is not needed. The AT, though, is not distribution free; therefore, the investigation of its validity over these situations is of interest. We also included the distribution-free procedures as a confirmation, to check the α s on the simulation and to identify the liberal and conservative tests by power.

As the underlying distribution changed from standard normal distribution to heavy tailed t-distribution with 1 degree of freedom, the empirical α for all tests are stable except the AT test. Both α s (.1217 and .0425) under t(1) are out of the 95% confidence intervals (.0941, .1059) and (.0457, .0543).

	N(0	(0, 1)	t(1)	LO)	t(5)	t(2)	t(1)
Meth.	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%
JT	.0982	.0469	.1049	.0501	.1047	.0490	.1093	.0521	.1017	.0489
SP	.0937	.0486	.1004	.0518	.0993	.0494	.1030	.0542	.0966	.0492
AT	.0961	.0473	.1001	.0501	.1004	.0507	.1095	.0534	.1217	.0425
HN	.0973	.0486	.1038	.0518	.1034	.0494	.1070	.0542	.1014	.0492

Table 3: Initial simulation on empirical α s.

2.1.2 Empirical Powers

For each simulation, n values (\mathbf{Z}) were generated from a specific distribution, and n_j of them (\mathbf{Z}_j) were randomly assigned to level j of the treatment. Suppose the center of the adjacent group is equally spaced by d. To specify the d value, the noncentrality parameter (δ) needs to first be found.

As in the null case, we set k = 5, $n_i = 5$, n = 25 and $\alpha = 0.05$. To determine d

select $p_0 = 0.15$. By solving the power expression

$$p_0 = P\left\{F(\delta, k-1, n-k) \ge F(1-\alpha, k-1, n-k)\right\},$$

we obtain the noncentrality parameter δ which is approximately 2.11.

Let X be the one-way ANOVA cell mean design matrix,

$$\boldsymbol{X}_{(n \times k)} = \begin{bmatrix} \boldsymbol{1}_{n_1} & \boldsymbol{0} & \dots & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{1}_{n_2} & \dots & \boldsymbol{0} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{0} & \boldsymbol{0} & \dots & \boldsymbol{1}_{n_k} \end{bmatrix},$$

and A be the matrix which consider group 1 as the reference group,

$$\boldsymbol{A}_{(k-1\times k)} = \begin{bmatrix} 1 & -1 & 0 & \dots & 0 \\ 1 & 0 & -1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 0 & 0 & \dots & -1 \end{bmatrix},$$

and

$$\boldsymbol{\beta} = \boldsymbol{\mu} = (\mu_1 \dots \mu_k)' = (\mu_1 \ \mu_1 + d \ \dots \ \mu_1 + (k-1) d)'.$$

We know that

$$\delta = \frac{1}{\sigma^2} \left(\boldsymbol{A} \boldsymbol{\mu} \right)^{\prime} \left[\boldsymbol{A} \left(\boldsymbol{X}^{\prime} \boldsymbol{X} \right)^{-1} \boldsymbol{A}^{\prime} \right]^{-1} \left(\boldsymbol{A} \boldsymbol{\mu} \right).$$

When $n_i = n_j$, after simplifying

$$\delta = \frac{n_i}{k\sigma^2} \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (\mu_i - \mu_j)^2$$

= $\frac{n_i}{k\sigma^2} \left[(k-1)1^2 + (k-2)2^2 + (k-3)3^2 + \dots + 1(k-1)^2 \right] d^2$

and setting $\sigma^2 = 1$ and $\mu_1 = 0$, we obtain $d = \sqrt{2.11/50}$.

Under the null hypothesis, all data are generated from the same distribution. While under the alternative hypothesis, all data are generated from the same distribution and then shifted in location. For example, the data for the jth group is

$$\boldsymbol{y}_j = \boldsymbol{Z}_j + \boldsymbol{b}_j, \text{ and } \boldsymbol{Y} = (\boldsymbol{y}_1 \ \boldsymbol{y}_2 \ \dots \ \boldsymbol{y}_k)',$$

where the b_j s represents the shifts of group j from group 1, and the values of b_j s are 0, $d, 2d, 3d, 4d \dots$ respectively, for $j = 1, \dots, k$.

Table 4 and Figure 2 show the empirical power and power curves when the errors have a normal distribution. For all tests, there is no significant difference among the empirical powers. Note that even though the errors are normally distributed, the AT test does not dominate the JT and HN tests.

Table 4: Initial simulation on empirical power under the Normal distribution.

Meth.	Empirical power				
	0	2.11	8.44		
JT	.0982	.5488	.9313		
SP	.0937	.5428	.9283		
AT	.0961	.5404	.9305		
HN	.0973	.5492	.9313		

Empirical Power Curves for 10% Tests



Figure 2: Plot of the initial study on power curves under a Normal distribution.

Empirical Power Curves for 10% Tests



Figure 3: Plot of the initial study on power curves under $t_{(10)}$.





Figure 4: Plot of the initial study on power curves under $t_{(8)}$.

Empirical Power Curves for 10% Tests



Figure 5: Plot of the initial study on power curves under $t_{(5)}$.



Empirical Power Curves for 10% Tests

Figure 6: Plot of the initial study on power curves under $t_{(3)}$.

Empirical Power Curves for 10% Tests



Figure 7: Plot of the initial study on power curves under $t_{(2)}$.



Empirical Power Curves for 10% Tests

Figure 8: Plot of the initial study on power curves under $t_{(1)}$.

Figures 3 to 8 show the empirical power curves under the t distributions with degrees of freedom 10, 8, 5, 3, 2, 1, respectively. As the tails of the distribution get heavier, the AT test becomes more and more instable. All distribution free tests are stable. Figure 8 shows that with t(1) error distribution, the JT test is more stable than the HN and SP tests, but there is no significant difference among them.

Meth.	Empirical power								
	0	2.11	8.44	18.99	33.76	52.75			
SP	.0995	.3227	.5920	.7783	.8660	.9149			
JT	.1051	.3324	.6107	.8034	.8959	.9424			
AT	.0952	.1517	.2126	.2893	.3715	.4536			
HN	.1034	.3292	.6013	.7830	.8702	.9176			

Table 5: Initial simulation on the empirical power under SCN distribution.



Empirical Power Curves for 10% Tests

Figure 9: Plot of the initial study on power curves under SCN distribution.

Table 5 and Figure 9 show the empirical power curves under the skewed contaminated normal distribution SCN(.25, 100, 1); the power of AT is quite low compared to other procedures. Statistically speaking, consider the standard two-sample, paired data, confidence interval for a difference in proportions. By using

$$\hat{p}_1 - \hat{p}_2 \pm Z_{\alpha/2} \sqrt{[\hat{p}_1 + \hat{p}_2 - (\hat{p}_1 - \hat{p}_2)^2]/n},$$

where *n* is the number of simulations, 10,000. This gives a confidence interval of the proportion difference at level α , to compare the methods. For our study, we used the last column empirical powers in table 5 with the JT, HN and SP tests having significantly higher power than the AT test with 95% confidence intervals (.4677, .5099), (.4429, .4851) and (.4402, .4824), respectively. The JT test does not have significantly higher power than the HN test with a 95% confidence interval (-.0019, .0515) and slightly higher power than the SP test with (.0008, .0542). The HN and SP tests are similar to each other with a 95% confidence interval (-.0238, .0292).

Based on all simulation results, the AT test is not as robust as other nonparametric approaches when the underlying error distribution has heavy tails. The first motivation of this thesis is to robustify the AT approach.

2.2 Robust Abelson-Tukey (RAT) Test

2.2.1 Rank-based Estimator

A general linear model is of the form

$$Y = \alpha + X\beta + e, \tag{2.1}$$

where \mathbf{Y} is the $n \times 1$ vector of responses, \mathbf{X} is the $n \times (k-1)$ design matrix, and \mathbf{e} is the $n \times 1$ vector of error terms. The least squares estimator minimizes the Euclidean distance between \mathbf{Y} and $\hat{\mathbf{Y}}$, the predicted value of \mathbf{Y} . For the robust Abelson-Tukey RAT procedure, a rank-based estimator, a different measure of distance based on the dispersion function of Jaeckel (Jaeckel, 1972; Jureckova, 1971) is used to substitute the LS estimator. The only assumption on the distribution of the errors required is that it is continuous.

As discussed in chapter 3 of Hettmansperger and McKean, the rank-based esti-

mator of $\boldsymbol{\beta}$ is defined as

$$\hat{\boldsymbol{\beta}} = \operatorname{Argmin} \| \boldsymbol{Y} - \boldsymbol{X} \boldsymbol{\beta} \|_{\omega},$$

where $\left\|\cdot\right\|_{\varphi}$ is a pseudo-norm defined as

$$\|u\|_{\varphi} = \sum_{i=1}^{n} a(R(u_i))u_i$$

where $R(u_i)$ denotes the rank of u_i , a(t) are scores such that $a(1) \leq \cdots \leq a(n)$ and

$$a(i) = \varphi\left(\frac{i}{n+1}\right),$$

where φ is a non-decreasing, square-integrable score function defined on the interval (0, 1). Assume without loss of generality that it is standardized, so that

$$\int \varphi(u) du = 0$$
 and $\int \varphi^2(u) du = 1$.

Kloke and McKean (2012) have developed a R package, Rfit, for rank-based (R) estimation and inference for linear models; see also (Kloke and McKean, 2014). In Rfit the default option is to use Wilcoxon (linear) scores. Furthermore, Rfit also includes a variaty of score functions, and user-defined score functions (Kloke and McKean, 2012) are easily added. The rank-based fit depends on a selection of score functions, which we discuss next.

2.2.2 General Rank Scores

To obtain efficient statistics, the density is assumed to have finite Fisher information. The Fisher information is given by $I(f) = \int_0^1 \varphi_f^2(u) du$, where

$$\varphi_f(u) = -\frac{f'\left(F^{-1}\left(u\right)\right)}{f\left(F^{-1}\left(u\right)\right)}$$
(2.2)

is the optimal score function.

When the errors come from different distributions, appropriate scores are recommended. Generally used scores include the following.

1. Wilcoxon scores are recommended when the errors come from a moderate tailed

distribution, and it is generated by the linear function $\varphi_R(u) = \sqrt{12}(u - 1/2)$. Wilcoxon scores are optimal if the error distribution is logistic. In Rfit, Wilcoxon scores can be set by scores=wscores.

- Normal scores are recommended when the errors come from a light-moderated tailed distribution. Normal scores are optimal for normal error distribution. In Rfit, normal scores can be set by scores=nscores.
- 3. There are four types of scores in the family of bent (Winsorized Wilcoxons) scores. They are optimal for error distribution with a logistic center and tails of exponential order(McKean et al., 1989).
 - (a) Bent1 scores are recommended when the errors come from a highly right skewed distribution, and the score function is defined in terms of 3 parameters. In Rfit, they can be set by scores=bentscores1.

$$\varphi(u) = \begin{cases} \frac{(s_3 - s_2)u}{s_1} & \text{if } u < s_1\\ s_3 & \text{else} \end{cases}$$

(b) Bent2 scores are recommended when the errors come from a light tailed distribution, and the score function is defined in terms of 5 parameters. In Rfit, they can be set by scores=bentscores2.

$$\varphi(u) = \begin{cases} \frac{(s_5 - s_3)u}{s_1} & \text{if } u < s_1\\ \frac{(s_4 - s_5)(u - s_2)}{1 - s_2} + s_5 & \text{if } u > s_2\\ s_5 & \text{else} \end{cases}$$

(c) Bent3 scores are recommended when the errors come from a highly left skewed distribution, and the score function is defined in terms of 3 parameters. In Rfit, they can be set by scores=bentscores3.

$$\varphi(u) = \begin{cases} s_2 & \text{if } u < s_1 \\ \frac{(s_3 - s_2)(u - s_1)}{1 - s_1} + s_2 & \text{else} \end{cases}$$

/

(d) Bent4 scores are recommended when the errors come from a moderately heavy tailed distribution, and the score function is defined in terms of 4 parameters. In Rfit, they can be set by scores=bentscores4.

$$\varphi(u) = \begin{cases} s_3 & \text{if } u < s_1 \\ s_4 & \text{if } u > s_2 \\ \frac{(s_4 - s_3)(u - s_1)}{s_2 - s_1} + s_3 & \text{else} \end{cases}$$

Plots of all these score functions are showed in Figure 10.

An appropriate bent score function for skewed distribution with a right heavy tail is as follows (Kloke and McKean, 2012):

$$\varphi(u) = \begin{cases} 4u - 1.5 & \text{if } u \le 0.5 \\ 0.5 & \text{if } u > 0.5 \end{cases}$$



Figure 10: Plots of score functions.

As shown in HM, the influence function of the rank-based estimation is bounded in the Y-space provided the score functions are bounded. Hence, all these score functions lead to robust fits in Y-space except for the normal scores fit. The normal scores estimator, however, is technically robust, since it has an unbounded influence function, but a positive breakdown point (Huber, 1981). The Wilcoxon procedures in linear models have the same efficiency properties as the Wilcoxon-Mann-Whitney procedures in location models. In particular, if the errors have a normal distribution this efficiency is 0.95. Generally, for distributions with heavier tails than normal, this efficiency is too large. For these reasons, the Wilcoxon scores are often used in practice. In this thesis, we often use Wilcoxon scores, but we also make use of other scores, especially the bent scores. For most of our discussion, the full model design is a one-way design. So there is no need for high breakdown estimates. A weighted version of the Wilcoxon can attain 50% breakdown in the X-space at the expense of a loss in efficiency (Chang et al., 1999).

2.2.3 The RAT Test

Let

$$\Delta = \boldsymbol{h}' \boldsymbol{\theta}_{i}$$

where $\boldsymbol{\theta}$ represents a $k \times 1$ vector of the location parameters of k treatment levels; \boldsymbol{h} is a $k \times 1$ vector of the AT maximin contrast coefficients. When $\hat{\boldsymbol{\theta}} = \bar{\boldsymbol{y}}$, where $\bar{\boldsymbol{y}}$ is a $k \times 1$ vector of the average of each treatment level, then the test statistic of the AT can be written as:

$$t = \frac{\boldsymbol{h}' \, \bar{\boldsymbol{y}}}{\sqrt{\operatorname{Var}(\boldsymbol{h}' \, \bar{\boldsymbol{y}})}}$$

Now, a rank-based estimator of Δ is obtained by using the rank-based linear model. Let \boldsymbol{W} be a $n \times k$ matrix which denotes the appropriate incidence matrix of 0s and 1s, and
note that the vector $\mathbf{1}_n$ is in the column space of \boldsymbol{W} , then

$$Y = W\theta + e$$

= $WEE^{-1}\theta + e$
= $X\beta + e$
= $\begin{bmatrix} 1 & X_1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta_1 \end{bmatrix} + e$
= $\alpha \mathbf{1} + X_1\beta_1 + e$,

where,

$$\boldsymbol{W}_{(n\times k)} = \begin{bmatrix} \mathbf{1}_{n_1} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{1}_{n_2} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{1}_{n_k} \end{bmatrix}$$
(2.3)
$$\boldsymbol{E}_{(k\times k)} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \\ 1 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & 1 \end{bmatrix}$$
(2.4)
$$\boldsymbol{X}_{\mathbf{1}(n\times (k-1))} = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{1}_{n_2} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \mathbf{1}_{n_3} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{1}_{n_k} \end{bmatrix}$$
(2.5)

and WE = X, $E^{-1}\theta = \beta$. Thus, the rank-based estimator of Δ is

$$\hat{\Delta} = \boldsymbol{h}' \hat{\boldsymbol{\theta}}$$

$$= \boldsymbol{h}' \boldsymbol{E} \hat{\boldsymbol{\beta}}$$

$$= \boldsymbol{h}' \begin{bmatrix} \mathbf{1} & \boldsymbol{E}_1 \end{bmatrix} \hat{\boldsymbol{\beta}}$$

$$= \begin{bmatrix} \mathbf{0} & \boldsymbol{h}' \boldsymbol{E}_1 \end{bmatrix} \hat{\boldsymbol{\beta}}$$

$$= \boldsymbol{h}' \boldsymbol{E}_1 \hat{\boldsymbol{\beta}}_1.$$

Under the assumption that the errors are iid with continuous probability density function f(t), it can be shown that the estimate $\hat{\beta}_1$ is consistent and asymptotically normal (Hettmansperger and McKean, 2011). The result can be summarized as follows:

$$\hat{\boldsymbol{\beta}}_{1} \dot{\sim} N(\boldsymbol{\beta}_{1}, \tau_{\varphi}^{2} (\boldsymbol{X}_{c}^{\prime} \boldsymbol{X}_{c})^{-1}), \qquad (2.6)$$

where X_c is centered X_1 , $X_c = (I - \frac{1}{n} \mathbf{11}') X_1$; where I is a $n \times n$ identity matrix and $\mathbf{1}' = (1, \ldots, 1)_{1 \times n}$; τ_{φ} is a scale parameter which depends on f, and the score function φ and is given by

$$\tau_{\varphi}^{-1} = \int \varphi(u)\varphi_f(u)du,$$
$$\varphi_f(u) = -\frac{f'(F^{-1}(u))}{f(F^{-1}(u))}.$$

Thus, the asymptotic distribution of $\hat{\Delta}$ is

$$\hat{\Delta} \sim N(\boldsymbol{h}' \boldsymbol{E}_1 \boldsymbol{\beta}_1, \tau_{\varphi}^2 \boldsymbol{h}' \boldsymbol{E}_1 (\boldsymbol{X}_c' \boldsymbol{X}_c)^{-1} \boldsymbol{E}_1' \boldsymbol{h}).$$
(2.7)

We denote this asymptotic variance by $\operatorname{Var}(\hat{\Delta})$.

Theorem 2.1. If $H_0: \theta_1 = \cdots = \theta_k$ is true, then $E(\hat{\Delta}) = 0$, as $n \to \infty$

$$RAT = \frac{\hat{\Delta}}{\sqrt{Var(\hat{\Delta})}} \xrightarrow{D} Z \sim N(0, 1).$$

For a level α test based on *RAT* for the hypothesis (1.3), suppose H_1 is true, then $\Delta = \mathbf{h}' \boldsymbol{\theta} > 0$. Since

$$\frac{1}{n} \boldsymbol{X}_{c}^{'} \boldsymbol{X}_{c} = O(1)$$
$$n(\boldsymbol{X}_{c}^{'} \boldsymbol{X}_{c})^{-1} = O(1),$$

and

$$(\boldsymbol{X}_{c}'\boldsymbol{X}_{c})^{-1} = \frac{1}{n}O(1),$$

therefore $\boldsymbol{h}' \boldsymbol{E}_1(\boldsymbol{X}'_c \boldsymbol{X}_c)^{-1} \boldsymbol{E}'_1 \boldsymbol{h} \to 0$ as $n \to \infty$. When *n* is sufficiently large, then we have

$$0 < z_{\alpha} \sqrt{\operatorname{Var}(\hat{\Delta})} < \Delta - \epsilon.$$

Hence,

$$P_{\Delta}[RAT \ge z_{\alpha}] = P_{\Delta}[\hat{\Delta} \ge z_{\alpha} \sqrt{\operatorname{Var}(\hat{\Delta})}]$$
$$\ge P_{\Delta}[\hat{\Delta} \ge \Delta - \epsilon]$$
$$\ge P_{\Delta}[\left|\hat{\Delta} - \Delta\right| \le \epsilon] \to 1,$$

thus the RAT test is consistent.

For a sequence of local alternatives $H_n: \theta_j = \theta_0 + n^{-1/2} c_j \theta^*$, where $\theta^* > 0$, j = 1, 2, ..., k. Let c be a vector of c_j , a set of constants, which specifies the pattern of the alternative. Suppose H_n is true, and F has a density f with $\int f^2(x) dx < \infty$, and suppose $\hat{\theta}_n$ is the rank based estimate of θ under H_n , and $\hat{\theta}_0$ is the rank based estimate of θ under H_0 thus,

$$\hat{\boldsymbol{ heta}}_n = \hat{\boldsymbol{ heta}}_0 + n^{-1/2} \theta^* \boldsymbol{c}$$

when H_n is true,

$$E(\hat{\Delta}) = E(\boldsymbol{h}'\hat{\boldsymbol{\theta}}_0 + n^{-1/2}\boldsymbol{\theta}^*\boldsymbol{h}'\boldsymbol{c}) = n^{-1/2}\boldsymbol{\theta}^*\boldsymbol{h}'\boldsymbol{c}$$

$$\operatorname{Var}(\hat{\Delta}) = \operatorname{Var}(\boldsymbol{h}'\hat{\boldsymbol{\theta}}_0) = \tau_{\varphi}^2 \boldsymbol{h}' \boldsymbol{E}_1(\boldsymbol{X}_c' \boldsymbol{X}_c)^{-1} \boldsymbol{E}_1' \boldsymbol{h}.$$

The asymptotic power of the test based on RAT is given by

$$P_{\Delta_n}[RAT \ge Z_{\alpha}] = P_{\Delta_n}[\hat{\Delta} \ge Z_{\alpha}\sqrt{\operatorname{Var}(\hat{\Delta})}]$$

= $P_{\Delta_n}[\frac{\hat{\Delta} - n^{-1/2}\theta^* \mathbf{h'c}}{\sqrt{\operatorname{Var}(\hat{\Delta})}} \ge Z_{\alpha} - \frac{n^{-1/2}\theta^* \mathbf{h'c}}{\sqrt{\operatorname{Var}(\hat{\Delta})}}]$
= $1 - \Phi(Z_{\alpha} - \theta^* c_{\varphi 0}),$

where $c_{\varphi 0}$ is the Pitman efficacy of the RAT test. This can be written as

$$c_{\varphi 0} = n^{-1/2} \boldsymbol{h}' \boldsymbol{c} / \sqrt{\operatorname{Var}(\hat{\Delta})} = n^{-1/2} \frac{\sum h_j c_j}{\tau_{\varphi} (\sum \frac{h_j^2}{n_j})^{1/2}}.$$

To obtain the asymptotic relative efficiency, an algebra form representing $\operatorname{Var}(\hat{\Delta})$ is given by

$$\operatorname{Var}(\hat{\Delta}) = \tau_{\varphi}^2 \sum_{j=1}^k \frac{h_j^2}{n_j},$$

since

$$\boldsymbol{h}'\boldsymbol{E}_1=[h_2\ h_3\ldots h_k],$$

and

$$(\boldsymbol{X}_{c}'\boldsymbol{X}_{c})_{(k-1)\times(k-1)}^{-1} = \begin{bmatrix} \frac{1}{n_{2}} + \frac{1}{n_{1}} & \frac{1}{n_{1}} & \dots & \frac{1}{n_{1}} \\ \frac{1}{n_{1}} & \frac{1}{n_{3}} + \frac{1}{n_{1}} & \dots & \frac{1}{n_{1}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n_{1}} & \frac{1}{n_{1}} & \dots & \frac{1}{n_{k}} + \frac{1}{n_{1}} \end{bmatrix}$$

$$\boldsymbol{h}' \boldsymbol{E}_{1} (\boldsymbol{X}_{c}' \boldsymbol{X}_{c})^{-1} (\boldsymbol{h}' \boldsymbol{E}_{1})' = \begin{pmatrix} \frac{h_{2}}{n_{2}} - \frac{h_{1}}{n_{1}} & \frac{h_{3}}{n_{3}} - \frac{h_{1}}{n_{1}} & \dots & \frac{h_{k}}{n_{k}} - \frac{h_{1}}{n_{1}} \end{pmatrix} \begin{pmatrix} h_{2} \\ h_{3} \\ \dots \\ h_{k} \end{pmatrix} = \sum_{j=1}^{k} \frac{h_{j}^{2}}{n_{j}}.$$

2.3 Simulation Results

In the previous sections it has been shown that the RAT test can use different score functions for different underlying distributions. The 10,000 simulation results in Table 6 show that RAT with Wilcoxon scores performs well, except that under t(1) it appears to be conservative. Because the t(1) distribution is symmetrical with heavy tails, the bent4 score would be more appropriate. We set $s_1 = 0.15$, $s_2 = 0.85$, $s_3 = -1$, $s_4 = 1$. For the bent4 score, the empirical levels of the corresponding RAT test, 0.0927 and 0.0487, were at nominal 10% and 5% levels respectively. Note that the RAT with the bent4 scores have empirical levels close to the nominal α under the t(1) distribution. Again, note the distribution freeness of the JT, SP and HN tests.

	N(0	N(0,1)		t(10)		5)	t(2)	t(1)	
Meth.	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%
SP	.0989	.0517	.1001	.0457	.1039	.0520	.1012	.0518	.0989	.0480
JT	.1036	.0509	.1016	.0461	.1079	.0503	.1046	.0522	.1024	.0477
AT	.1013	.0495	.0994	.0479	.1038	.0512	.1065	.0504	.1191	.0414
RAT	.1026	.0512	.0982	.0504	.1010	.0493	.0921	.0469	.0729	.0355
RATb4	.1168	.0638	.1128	.0613	.1124	.0625	.1078	.0591	.0927	.0487
HN	.1038	.0517	.1034	.0457	.1078	.0520	.1050	.0518	.1023	.0480

Table 6: Empirical α s at nominal levels.

Table 7 shows the empirical powers of all five methods with different underlying distributions. The JT and HN tests have the highest empirical power at each simulation. Their empirical powers are not significantly different.

	N(0	N(0,1)		t(10)		5)	t(2)	t(1)	
Meth.	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%
SP	.5433	.3966	.5108	.3629	.4638	.3344	.3948	.2598	.3094	.1896
JT	.5519	.3909	.5218	.3583	.4728	.3287	.4048	.2525	.3173	.1926
AT	.5401	.3943	.4957	.3453	.4356	.3000	.2925	.1770	.1841	.0832
RAT	.5168	.3640	.4844	.3332	.4384	.3020	.3521	.2174	.2196	.1205
RATb4	.5248	.3803	.4963	.3538	.4525	.3204	.3726	.2450	.2444	.1448
HN	.5516	.3966	.5184	.3629	.4715	.3344	.4027	.2598	.3163	.1896

Table 7: Empirical power at nominal levels.

Table 8 and Figure 11 show the empirical power of five methods under normal distribution. There is no significant difference.

Table 9 and Figure 12 show the empirical power of all five methods under the skewed contaminated normal distribution (SCN(.25, 100, 1)). If we use the last column of

the empirical power of table 9 to compute a 95% confidence interval of the proportion difference, we find the RAT method has higher power than AT methods under the skewed contaminated normal distribution with (.3135, .3603), and especially in the RAT with bent1 score method, with (.3402, .3872).

Meth.]	Empirical power								
	0	2.11	8.44	18.99						
SP	.1006	.5433	.9341	.9980						
JT	.1030	.5519	.9356	.9978						
AT	.1012	.5401	.9354	.9978						
RAT	.1050	.5168	.9143	.9964						
RATb1	.1174	.5248	.9060	.9951						
HN	.1035	.5516	.9374	.9980						

Table 8: Empirical power levels under the Normal distribution.

In figure 12, the power curve of the RAT method is below the JT, HN, and SP methods, but the power curve of the RAT with bent1 score method crosses the curves of the HN and SP methods, and falls right below the JT curve. There are significant differences between RAT and AT, and between RATb1 and AT. The RAT and RATb1 tests are more powerful than the AT test. The 95% confidence interval of RATb1 and JT shows there is no significant difference between them with (-.0397, .0149).

Empirical Power Curves for 10% Tests



Figure 11: Plot of power curves under a Normal distribution.

Meth.		Empirical power											
	0	2.11	8.44	18.99	33.76	52.75	75.96	103.39					
SP	.1079	.3233	.5953	.7763	.8657	.9150	.9395	.9548					
JT	.1109	.3317	.6151	.8011	.8994	.9420	.9660	.9791					
AT	.1045	.1531	.2123	.2886	.3717	.4540	.5299	.6030					
RAT	.0714	.1780	.3671	.5867	.7566	.8603	.9132	.9399					
RATb1	.0910	.2708	.5394	.7575	.8729	.9285	.9584	.9667					
HN	.1112	.3314	.6035	.7818	.8719	.9179	.9416	.9563					

Table 9: Empirical power levels under the SCN distribution.

Empirical Power Curves for 10% Tests



Figure 12: Plot of power curves under the SCN distribution

Based on all the simulation results, we can conclude that the RAT test is robust, it allows any scores, and it did better than the AT test. Under the skewed contaminated normal distribution (SCN(.25, 1, 100)) the RAT with bent1 scores test does not outperform the JT test.

Terpstra (1952) and Jonckheere (1954) proposed the test based on pairwise Mann-Whitney-Wilcoxon statistics, while RAT only uses partial shifts based on the number of groups (k) as even or odd. Such as when k = 5, there are 10 U_{ij} s in the JT statistic, while only 2 Δ_{ij} s exist in the RAT statistic. In the JT statistic:

$$J = \sum_{i < j} \sum_{i < j} U_{ij}$$

= $U_{12} + U_{13} + U_{14} + U_{15} + U_{23} + U_{24} + U_{25} + U_{34} + U_{35} + U_{45}.$

In the RAT statistic:

$$\Delta = c'\theta$$

= -0.89\theta_1 - 0.2\theta_2 + 0.2\theta_4 + 0.89\theta_5
= 0.89(\theta_5 - \theta_1) + 0.2(\theta_4 - \theta_2)
= 0.89\Delta_{51} + 0.2\Delta_{42}.

Thus, JT uses more information than RAT, and the pairwise statistics provide more direct information on the sources of statistical significance. In light of this, we consider the Hettmansperger-Norton optimal weights to be a better choice.

CHAPTER 3

NEW METHODS

Recall that in this k-sample location problem, the response Y_{ij} follows the linear model,

$$Y_{ij} = \theta_j + \epsilon_{ij}, \ j = 1, \dots, \ k, \ i = 1, \dots, \ n_j.$$

 Y_{ij} is the *ith* observation on response variable for the *jth* treatment level; θ_j is the *jth* location parameter; ϵ_{ij} are independent and identically distributed (*iid*) with cdf F(x) and pdf f(x). The null hypothesis of interest is that there is no differences in locations among k populations, or, there is no difference in treatment effects (i.e. 1.2), and the alternative hypotheses of interest follow a pattern (i.e. (1.3), (1.4) or (1.5)).

In this chapter, three new methods, the robust Abelson-Tukey with HN weights (RAThn) test, the Robust Shao-McKean (RSM) test and the RSM with least square estimates (RSM_{LS}) test, are presented. All three methods have the same weights, the HN weights, and the same form of a test statistic, but have different estimates of the shifts.

The test statistics used in this chapter have the form:

$$T = \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) \hat{\Delta}_{ji}, \qquad (3.1)$$

where $a_j = \lambda_j (c_j - \bar{c}_w)$, c_j specifics the pattern of the alternative, and $\bar{c}_w = \sum \lambda_j c_j$, $\lambda_j = \frac{n_j}{n}$ and $\sum_{j=1}^k a_j = 0$. $\hat{\Delta}_{ji}$ is an estimate which measures the shift from group j to group i. The robust Abelson-Tukey with HN weights (RAThn) test uses the rank-based regression estimate; the Robust Shao-McKean (RSM) test uses a pseudo-median estimate, the Hodges-Lehmann estimate; the RSM with least square estimates (RSM_{LS}) test uses the least square estimate to estimate the shift $\hat{\Delta}_{ji}$.

3.1 Robust Abelson-Tukey with HN Weights (RAThn) Test

Recall in section 2.2 that a rank-based estimate of β (the shifts of other groups from the first group) is defined as

$$\hat{\boldsymbol{\beta}} = \operatorname{Argmin} \| \boldsymbol{Y} - \boldsymbol{X} \boldsymbol{\beta} \|_{\omega},$$

and under the assumption that the errors are iid with continuous probability density function, the estimate $\hat{\beta}$ is consistent and asymptotically normal, which is shown as 2.6. Even though it does not give the rank-based estimates of the pairwise shifts directly, but the following lemma shows that the test statistic can be represented as a linear combination of the shifts of all groups from the first group, $\hat{\Delta}_{i1}$, $i = 1, 2, \ldots k$.

Lemma 1. Suppose $\Delta_{ji} = \Delta_{j1} - \Delta_{i1}$, where $\sum_{j=1}^{k} a_j = 0$ and $\Delta_{11} = 0$. Then

$$\sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) \Delta_{ji} = k \sum_{i=1}^{k} a_i \Delta_{i1}.$$

Proof:

$$\sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) \Delta_{ji} = \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) (\Delta_{j1} - \Delta_{i1})$$
$$= \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} a_j \Delta_{j1} - \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} a_i \Delta_{j1}$$
$$- \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} a_j \Delta_{i1} + \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} a_i \Delta_{i1}$$

$$= \sum_{i=1}^{k} (i-1)a_{i}\Delta_{i1} + \sum_{i=1}^{k-1} (k-i)a_{i}\Delta_{i1}$$

$$-\sum_{i=2}^{k} \sum_{j < i} a_{j}\Delta_{i1} - \sum_{i=1}^{k-1} \sum_{j > i} a_{j}\Delta_{i1}$$

$$= \sum_{i=1}^{k-1} (i-1)a_{i}\Delta_{i1} + (k-1)a_{k}\Delta_{k1} + \sum_{i=1}^{k-1} (k-i)a_{i}\Delta_{i1} - \sum_{i=1}^{k} \sum_{j \neq i} a_{j}\Delta_{i1}$$

$$= \sum_{i=1}^{k-1} (k-i+i-1)a_{i}\Delta_{i1} + (k-1)a_{k}\Delta_{k1} - \sum_{i=1}^{k} \Delta_{i1}\sum_{j \neq i} a_{j}$$

$$= \sum_{i=1}^{k-1} (k-1)a_{i}\Delta_{i1} + (k-1)a_{k}\Delta_{k1} + \sum_{i=1}^{k} \Delta_{i1}a_{i}$$

$$= (k-1)\sum_{i=1}^{k} a_{i}\Delta_{i1} + \sum_{i=1}^{k} \Delta_{i1}a_{i}$$

$$= k\sum_{i=1}^{k} a_{i}\Delta_{i1}$$

$$= k\sum_{i=1}^{k} a_{i}\Delta_{i1}.$$

$$(\sum_{j=1}^k a_j = 0 \Rightarrow \sum_{j \neq i} a_j = -a_i).$$

3.1.1 The RAThn Test Statistic

The RAThn test statistic is

$$T = k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_1, \tag{3.2}$$

where $\hat{\boldsymbol{\beta}}_1$ is a vector of the estimates of the shifts from group *i* to group 1, $\underline{\boldsymbol{a}} = (a_2, a_3, \ldots, a_k)'$, where $\sum_{i=1}^k a_i = 0$. Based on (2.6) the asymptotically distribution of *T* is

$$T \dot{\sim} N(k \boldsymbol{a}' \boldsymbol{\beta}_1, k^2 \tau_{\varphi}^2 \boldsymbol{a}' (\boldsymbol{X}_c' \boldsymbol{X}_c)^{-1} \boldsymbol{a}).$$
(3.3)

Theorem 3.1. If H_0 : $\theta_1 = \cdots = \theta_k$ is true, then E(T) = 0. and Var(T) =

 $k^2 \hat{\tau}_{\varphi}^2 a' (X_c' X_c)^{-1} a$. Then as $n \to \infty$, where $n = \sum_{i=1}^k n_i$,

$$T^* = \frac{k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_1}{\sqrt{Var(T)}} \xrightarrow{D} Z \sim N(0, 1).$$

The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3)

if $T^* \geq z_{\alpha}$; otherwise do not reject.

3.1.2 Consistency and Asymptotic Power

Suppose H_1 is true, then $T = ka'\beta_1 > 0$.

Since

$$\frac{1}{n} \boldsymbol{X}_{c}^{'} \boldsymbol{X}_{c} = O(1)$$
$$n(\boldsymbol{X}_{c}^{'} \boldsymbol{X}_{c})^{-1} = O(1),$$

and

$$(\boldsymbol{X}_{c}'\boldsymbol{X}_{c})^{-1} = \frac{1}{n}O(1),$$

thus, $\boldsymbol{a}'(\boldsymbol{X}_c'\boldsymbol{X}_c)^{-1}\boldsymbol{a} \to 0 \text{ as } n \to \infty.$

If n is sufficiently large, then we have

$$0 < z_{\alpha} \sqrt{\operatorname{Var}(T)} < ka' \beta_1 - \epsilon.$$

Hence,

$$P[T^* \ge z_{\alpha}] = P\left[T \ge z_{\alpha}\sqrt{\operatorname{Var}(T)}\right]$$
$$\ge P\left[ka'\hat{\beta}_1 \ge ka'\beta_1 - \epsilon\right]$$
$$\ge P\left[\left|\hat{\beta}_1 - \beta_1\right| \le \epsilon\right] \to 1.$$

Thus, RAThn is a consistent test.

Under a sequence of alternatives H_n : $\theta_j = \theta_0 + \theta c_j n^{-1/2}$, where $\theta > 0$, j =

 $1, \ldots, k$, and F has a density f with $\int f^2(x) dx < \infty$,

$$\Delta_{i1} = (c_i - c_1)\theta n^{-1/2}.$$

Let D denote the vector of Δ_{i1} s, and C denote the vector of $c_i - c_1$, so

$$\boldsymbol{D}_n = \boldsymbol{C}\theta n^{-1/2}.$$

When H_n is true,

$$\mathbf{E}(T) = k \boldsymbol{a}' \boldsymbol{C} \boldsymbol{\theta} n^{-1/2}$$

$$\operatorname{Var}(T) = k^{2} \tau_{\varphi}^{2} \boldsymbol{a}' (\boldsymbol{X}_{c}' \boldsymbol{X}_{c})^{-1} \boldsymbol{a}.$$

The asymptotic power of the test based on T is given by

$$P_{D_n}[T^* \ge Z_\alpha] = P_{D_n}[T \ge Z_\alpha \sqrt{\operatorname{Var}(T)}]$$

= $P_{D_n}[\frac{T - ka'D_n}{\sqrt{\operatorname{Var}(T)}} \ge Z_\alpha - \frac{ka'D_n}{\sqrt{\operatorname{Var}(T)}}]$
= $1 - \Phi(Z_\alpha - \theta c_{\varphi 1}),$

where $c_{\varphi 1}$ is the Pitman efficacy of the RAThn test, and it can be written as

$$c_{\varphi 1} = a' C \frac{k n^{-1/2}}{\sqrt{k^2 \tau_{\varphi}^2}} (a' (X'_c X_c)^{-1} a)^{-1/2}$$

$$= \frac{1}{\sqrt{n} \tau_{\varphi}} a' C (a' (X'_c X_c)^{-1} a)^{-1/2}$$

$$= \frac{\sum_{i=1}^k a_i c_i}{\tau_{\varphi} (\sum_{i=1}^k \frac{a_i^2}{\lambda_i})^{1/2}}.$$

3.2 Robust Shao-McKean (RSM) Test

Hettmansperger-Norton (1987) used a linear combination of pairwise comparisons based on the MWW statistic $U_{ij} = \sum_{m=1}^{n_i} \sum_{l=1}^{n_j} \phi(Y_{mi}, Y_{lj}), (1 \le i < j \le k),$ where $\phi(a, b) = 1$, if a < b, 0 otherwise. Our effect-size test is based on a linear combination of the estimates $\hat{\Delta}_{ij}$.

Suppose that Y_{ij} is distributed with continuous cdf $F(y - \theta_j)$ for $j = 1, \ldots, k$ and $i = 1, \ldots, n_j$ and the observations are mutually independent. We want to test $H_0: \theta_1 = \cdots = \theta_k$ versus $H_A: \theta_j = \theta_0 + \theta_{cj}, (\theta > 0; j = 1, \ldots, k)$, where c_1, \ldots, c_k is a given set of constants that specifies the pattern of the alternative. The experiment may specify the pattern and spacings based on previous studies and/or theoretical formulations. Equally spaced constants are recommended unless there are indications to the contrary.

3.2.1 The RSM Test Statistic

For any pattern with a known peak, the RSM test statistic is

$$RSM = \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) \hat{\Delta}_{ij} = \sum_{i < j} \sum_{i < j} a_{ij} \hat{\Delta}_{ij}.$$
 (3.4)

The Hodges-Lehmann estimate of shift for Δ_{ij} is

$$\hat{\Delta}_{ij} = med\left\{Y_{lj} - Y_{l'i}\right\}, \ 1 \le l \le n_j, \ 1 \le l' \le n_i.$$
(3.5)

Under the null hypothesis E(RSM) = 0, and

$$Var(RSM) = \sum_{i < j} \sum_{i' < j'} a_{ij} a_{i'j'} Cov(\hat{\Delta}_{ij}, \hat{\Delta}_{i'j'}).$$

$$(3.6)$$

From (HM2.5.30), (HM2.5.20) and (HM2.5.5) we have the asymptotic representation

$$\sqrt{n_{ij}}\hat{\Delta}_{ij} = \frac{\tau_{\varphi}}{\sqrt{n_{ij}}\lambda_i^{ij}\lambda_j^{ij}}T_{\varphi}(0) + o_p(1), \qquad (3.7)$$

where

$$T_{\varphi}(0) = \sum_{i'=1}^{n_{ij}} (b_{i'} - \bar{b}_{ij})\varphi\left[F(Z_{i'})\right], \qquad (3.8)$$

where $n_{ij} = n_i + n_j$, $b_{i'} = 1$ if $n_i + 1 \le i' \le n_{ij}$; $b_{i'} = 0$ otherwise, $\lambda_i^{ij} = n_i/n_{ij}$, $\bar{b}_{ij} = \sum b_i/n_{ij} = n_j/n_{ij}$, $(1 - \bar{b}_{ij}) = n_i/n_{ij}$ and

$$Z_{i'} = \Delta_{ij} b_{i'} + e_{i'}, \ 1 \ \le \ i' \ \le \ n_{ij},$$

and where $e_1, \ldots e_{n_{ij}}$ are *iid* with distribution function F(x). And

$$\int_{0}^{1} \varphi(u) d_{u} = 0, \text{ and } \int_{0}^{1} \varphi^{2}(u) d_{u} = 1.$$
(3.9)

Thus,

$$\begin{split} Cov(\hat{\Delta}_{ij},\hat{\Delta}_{i'j'}) &= \frac{\tau_{\varphi}^2}{n_{ij}n_{i'j'}\lambda_{i}^{ij}\lambda_{jj'}^{ij}\lambda_{i'}^{i'j'}}\sum_{m=1}^{n_{ij}}\sum_{l=1}^{n_{ij}'}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})\\ &\quad cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'})))\\ &= \kappa\sum_{m=1}^{n_i}\sum_{l=1}^{n_i'}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'})))\\ &\quad +\kappa\sum_{m=1}^{n_i}\sum_{l=i'+1}^{n_{i'j'}}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'})))\\ &\quad +\kappa\sum_{m=n_i+1}^{n_{ij}}\sum_{l=1}^{n_{i'}}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'})))\\ &\quad +\kappa\sum_{m=n_i+1}^{n_{ij}}\sum_{l=n_{i'}+1}^{n_{i'j'}}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'}))))\\ &\quad +\kappa\sum_{m=n_i+1}^{n_{ij}}\sum_{l=n_{i'}+1}^{n_{i'j'}}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'}))))\\ &\quad +\kappa\sum_{m=n_i+1}^{n_{ij}}\sum_{l=n_{i'}+1}^{n_{i'j'}}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'}))))\\ &\quad +\kappa\sum_{m=n_i+1}^{n_{ij}}\sum_{l=n_{i'}+1}^{n_{i'j'}}(b_m^{ij}-\bar{b}_{ij})(b_l^{i'j'}-\bar{b}_{i'j'})cov(\varphi(F(Z_m^{ij})),\varphi(F(Z_l^{i'j'})))), \end{split}$$

where $\kappa = \frac{\tau_{\varphi}^2}{n_{ij}n_{i'j'}\lambda_i^{ij}\lambda_j^{ij}\lambda_j^{i'j'}\lambda_{j'}^{i'j'}}$.

• $cov(\hat{\Delta}_{ij},\hat{\Delta}_{i'j'})=0$, if there are no ties among i,j,i',j';

• if $i = i', cov(\hat{\Delta}_{ij}, \hat{\Delta}_{i'j'}) = \kappa \sum_{m=1}^{n_i} \sum_{l=1}^{n_{i'}} (b_m^{ij} - \bar{b}_{ij}) (b_l^{i'j'} - \bar{b}_{i'j'}) = \frac{\tau_{\varphi}^2}{n_i};$

• if
$$j = j', cov(\hat{\Delta}_{ij}, \hat{\Delta}_{i'j'}) = \kappa \sum_{m=n_i+1}^{n_{ij}} \sum_{l=i'+1}^{n_{i'j'}} (b_m^{ij} - \bar{b}_{ij})(b_l^{i'j'} - \bar{b}_{i'j'}) = \frac{\tau_{\varphi}^2}{n_j};$$

• if
$$i = j', cov(\hat{\Delta}_{ij}, \hat{\Delta}_{i'j'}) = \kappa \sum_{m=1}^{n_i} \sum_{l=i'+1}^{n_{i'j'}} (b_m^{ij} - \bar{b}_{ij}) (b_l^{i'j'} - \bar{b}_{i'j'}) = -\frac{\tau_{\varphi}^2}{n_i};$$

• if
$$j = i', cov(\hat{\Delta}_{ij}, \hat{\Delta}_{i'j'}) = \kappa \sum_{m=n_i+1}^{n_{ij}} \sum_{l=1}^{n_{i'}} (b_m^{ij} - \bar{b}_{ij}) (b_l^{i'j'} - \bar{b}_{i'j'}) = -\frac{\tau_{\varphi}^2}{n_j};$$

Also, the asymptotic variance of $\hat{\Delta}_{ij}$ is

$$\begin{aligned} AV(\hat{\Delta}_{ij}) &= \frac{\tau_{\varphi}^2}{n_{ij}^2 \lambda_i^2 \lambda_j^2} \sum_{i'=1}^{n_{ij}} (b_{i'} - \bar{b}_{ij})^2 Var\left[\varphi(F(Z_{i'}))\right] \\ &= \frac{\tau_{\varphi}^2}{n_{ij}^2 \lambda_i^2 \lambda_j^2} \sum_{i'=1}^{n_{ij}} (b_{i'} - \bar{b}_{ij})^2 \\ &= \frac{\tau_{\varphi}^2 n_{ij}^2}{n_i^2 n_j^2} \left[n_i (0 - \bar{b}_{ij})^2 + n_j (1 - \bar{b}_{ij})^2 \right] \\ &= \frac{\tau_{\varphi}^2 n_{ij}^2}{n_i^2 n_j^2} \left[\frac{n_i n_j^2}{n_{ij}^2} + \frac{n_i^2 n_j}{n_{ij}^2} \right] \\ &= \frac{\tau_{\varphi}^2}{n_i^2 n_j^2} n_i n_j (n_i + n_j) \\ &= \frac{\tau_{\varphi}^2}{n_i n_j} n_{ij} \\ &= \tau_{\varphi}^2 (\frac{1}{n_i} + \frac{1}{n_j}). \end{aligned}$$

From (3.4) we have

$$RSM = \sum_{i < j} \sum_{i < j} (a_i - a_j) \hat{\Delta}_{ij}.$$

If we let

$$RSM = \sum \sum_{i < j} v_{ij}$$

Therefore,

$$Var(RSM) = \sum_{i < j} \sum_{i < j} Var(v_{ij}) + \sum_{i < j} \sum_{i < j} \sum_{i' = 1, i' \neq i}^{j-1} Cov(v_{ij}, v_{i'j}) + \sum_{j' = i+1, j' \neq j}^{k} Cov(v_{ij}, v_{ij'}) + \sum_{i' = j+1, i' \neq i}^{k} Cov(v_{ij}, v_{ji'}) + \sum_{j' = 1, j' \neq j}^{i-1} Cov(v_{ij}, v_{j'j})]$$

continuously,

$$\begin{aligned} Var(RSM) &= \sum_{i < j} \sum_{i < j} (a_j - a_i)^2 \tau_{\varphi}^2 (\frac{1}{n_i} + \frac{1}{n_j}) \\ &+ \sum_{i < j} \sum_{i < j} \sum_{i'=1, i' \neq i}^{j-1} (a_j - a_i)(a_j - a_{i'}) \frac{\tau_{\varphi}^2}{n_j} \\ &+ \sum_{j'=i+1, j' \neq j}^k (a_j - a_i)(a_{j'} - a_i) \frac{\tau_{\varphi}^2}{n_i} \\ &- \sum_{i'=j+1, i' \neq i}^k (a_j - a_i)(a_i - a_j) \frac{\tau_{\varphi}^2}{n_j} \\ &- \sum_{j'=1, j' \neq j}^{i-1} (a_j - a_i)(a_i - a_{j'}) \frac{\tau_{\varphi}^2}{n_i}] \\ &= \tau_{\varphi}^2 \sum_{i < j} \sum_{i < j} (a_j - a_i) \sum_{i'=1, i' \neq i, j}^k (a_j - a_{i'}) \frac{1}{n_j} \\ &+ \tau_{\varphi}^2 \sum_{i < j} \sum_{i < j} (a_j - a_i) \sum_{i'=1, i' \neq i, j}^k (a_j - a_{i'}) \frac{1}{n_j} \end{aligned}$$

Since $\sum_{i'=1,i'\neq i,j}^{k} a_j = (k-2)a_j$, and $\sum_{i'=1,i'\neq i,j}^{k} a_{i'} = -(a_i+a_j)$, $(\sum a_i = 0)$, Var(RSM) simplifies to

$$\begin{aligned} Var(RSM) &= \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})^{2} (\frac{1}{n_{i}} + \frac{1}{n_{j}}) \\ &+ \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i}) [\frac{1}{n_{j}} [(k - 2)a_{j} + a_{i} + a_{j}] - \frac{1}{n_{i}} [(k - 2)a_{i} + a_{i} + a_{j}]] \\ &= \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})^{2} (\frac{1}{n_{i}} + \frac{1}{n_{j}}) \\ &+ \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i}) [k(\frac{a_{j}}{n_{j}} - \frac{a_{i}}{n_{i}}) - (a_{j} - a_{i})(\frac{1}{n_{i}} + \frac{1}{n_{j}})] \\ &= \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})^{2} (\frac{1}{n_{i}} + \frac{1}{n_{j}}) \\ &+ k\tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})(\frac{a_{j}}{n_{j}} - \frac{a_{i}}{n_{i}}) - \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})^{2} (\frac{1}{n_{i}} + \frac{1}{n_{j}}) \\ &= k\tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})(\frac{a_{j}}{n_{j}} - \frac{a_{i}}{n_{i}}) \\ &= k\tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})(\frac{a_{j}}{n_{j}} - \frac{a_{i}}{n_{i}}) \\ &= k\tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})(\frac{a_{j}}{n_{j}} - \frac{a_{j}}{n_{j}}a_{i} - a_{j}\frac{a_{i}}{n_{i}} + \frac{a_{i}^{2}}{n_{i}}) \\ &= k\tau_{\varphi}^{2} \sum_{j = 2} \sum_{i < j} (j - 1)\frac{a_{j}^{2}}{n_{j}} - \frac{a_{j}}{n_{j}}\sum_{i = 1}^{j - 1} a_{i} - a_{j}\sum_{i = 1}^{j - 1} \frac{a_{i}}{n_{i}} + \sum_{i = 1}^{j - 1} \frac{a_{i}^{2}}{n_{i}} \end{bmatrix}$$

$$\sum_{j=2}^{k} \frac{a_j}{n_j} \sum_{i=1}^{j-1} a_i = \sum_{j=2}^{k-1} \frac{a_j}{n_j} \sum_{i=1}^{j-1} a_i + \frac{a_k}{n_k} \sum_{i=1}^{k-1} a_i$$
$$= \sum_{j=2}^{k-1} \frac{a_j}{n_j} \sum_{i=1}^{j-1} a_i - \frac{a_k^2}{n_k}$$

$$\sum_{j=2}^{k} a_j \sum_{i=1}^{j-1} \frac{a_i}{n_i} = \sum_{j=1}^{k-1} \frac{a_j}{n_j} \sum_{i=j+1}^{k} a_i$$
$$= \frac{a_1}{n_1} \sum_{i=2}^{k} a_i + \sum_{j=2}^{k-1} \frac{a_j}{n_j} \sum_{i=j+1}^{k} a_i$$
$$= -\frac{a_1^2}{n_1} + \sum_{j=2}^{k-1} \frac{a_j}{n_j} \sum_{i=j+1}^{k} a_i$$

$$\begin{split} \sum_{j=2}^{k-1} \frac{a_j}{n_j} \sum_{i=1}^{j-1} a_i - \frac{a_k^2}{n_k} - \frac{a_1^2}{n_1} + \sum_{j=2}^{k-1} \frac{a_j}{n_j} \sum_{i=j+1}^k a_i &= -\frac{a_k^2}{n_k} - \frac{a_1^2}{n_1} + \sum_{j=2}^{k-1} \frac{a_j}{n_j} \left[\sum_{i=1}^{j-1} a_i + \sum_{i=j+1}^k a_i \right] \\ &= -\frac{a_k^2}{n_k} - \frac{a_1^2}{n_1} - \sum_{j=2}^{k-1} \frac{a_j^2}{n_j} \\ &= -\sum_{j=1}^k \frac{a_j^2}{n_j}. \end{split}$$

Thus,

$$\begin{aligned} Var(RSM) &= k\tau_{\varphi}^{2} \left[\sum_{j=2}^{k} (j-1) \frac{a_{j}^{2}}{n_{j}} + \sum_{j=1}^{k} \frac{a_{j}^{2}}{n_{j}} + \sum_{j=2}^{k} \sum_{i=1}^{j-1} \frac{a_{i}^{2}}{n_{i}} \right] \\ &= k\tau_{\varphi}^{2} \left[\sum_{j=1}^{k} (j-1) \frac{a_{j}^{2}}{n_{j}} + \sum_{j=1}^{k} \frac{a_{j}^{2}}{n_{j}} + \sum_{j=1}^{k-1} (k-j) \frac{a_{j}^{2}}{n_{j}} \right] \\ &= k\tau_{\varphi}^{2} \left[\sum_{j=1}^{k} (j-1) \frac{a_{j}^{2}}{n_{j}} + \sum_{j=1}^{k} \frac{a_{j}^{2}}{n_{j}} + \sum_{j=1}^{k} (k-j) \frac{a_{j}^{2}}{n_{j}} \right] \\ &= k^{2} \tau_{\varphi}^{2} \sum_{j=1}^{k} \frac{a_{j}^{2}}{n_{j}}. \end{aligned}$$

For asymptotic theory, consider the asymptotic representation of $\hat{\Delta}_{ij}$; i.e.

$$\begin{split} \sqrt{n_{ij}}\hat{\Delta}_{ij} &= \frac{\tau_{\varphi}}{\sqrt{n_{ij}}\lambda_{i}^{ij}\lambda_{j}^{ij}} \sum_{m=1}^{n_{ij}} (c_{m} - \bar{c}_{ij})\varphi\left[F(Z_{m})\right] + o_{p}(1) \\ &= \frac{\tau_{\varphi}}{\sqrt{n_{ij}}\lambda_{i}^{ij}\lambda_{j}^{ij}} \left\{ \sum_{m=1}^{n_{ij}} c_{m}\varphi\left[F(Z_{m})\right] - \bar{c}_{ij}\sum_{m=1}^{n_{ij}}\varphi\left[F(Z_{m})\right] \right\} + o_{p}(1) \\ &= \frac{\tau_{\varphi}}{\sqrt{n_{ij}}\lambda_{i}^{ij}\lambda_{j}^{ij}} \left\{ \sum_{m=n_{i}+1}^{n_{ij}}\varphi\left[F(Z_{m})\right] - \frac{n_{j}}{n_{ij}}\sum_{m=1}^{n_{ij}}\varphi\left[F(Z_{m})\right] \right\} + o_{p}(1) \\ &= \frac{\tau_{\varphi}}{\sqrt{n_{ij}}\lambda_{i}^{ij}\lambda_{j}^{ij}} \left\{ \frac{n_{i}}{n_{ij}}\sum_{m=n_{i}+1}^{n_{ij}}\varphi\left[F(Z_{m})\right] - \frac{n_{j}}{n_{ij}}\sum_{m=1}^{n_{i}}\varphi\left[F(Z_{m})\right] \right\} + o_{p}(1) \\ &= \frac{\tau_{\varphi}}{\sqrt{n_{ij}}\lambda_{i}^{ij}\lambda_{j}^{ij}} \left\{ \lambda_{i}^{ij}\sum_{m=n_{i}+1}^{n_{ij}}\varphi\left[F(Z_{m})\right] - \lambda_{j}^{ij}\sum_{m=1}^{n_{i}}\varphi\left[F(Z_{m})\right] \right\} + o_{p}(1) \\ &= \frac{\tau_{\varphi}}{\sqrt{n_{ij}}} \left\{ \frac{1}{\lambda_{j}^{ij}}\sum_{m=n_{i}+1}^{n_{ij}}\varphi\left[F(Z_{m})\right] - \frac{1}{\lambda_{i}^{ij}}\sum_{m=1}^{n_{i}}\varphi\left[F(Z_{m})\right] \right\} + o_{p}(1), \end{split}$$

$$\begin{aligned} \hat{\Delta}_{ij} &= \frac{\tau_{\varphi}}{n_{ij}} \left\{ \frac{1}{\lambda_{j}^{ij}} \sum_{m=n_{i}+1}^{n_{ij}} \varphi\left[F(Z_{m})\right] - \frac{1}{\lambda_{i}^{ij}} \sum_{m=1}^{n_{i}} \varphi\left[F(Z_{m})\right] \right\} + o_{p}(\frac{1}{\sqrt{n_{ij}}}) \\ &= \tau_{\varphi} \left\{ \frac{1}{n_{j}} \sum_{m=n_{i}+1}^{n_{ij}} \varphi\left[F(Z_{m})\right] - \frac{1}{n_{i}} \sum_{m=1}^{n_{i}} \varphi\left[F(Z_{m})\right] \right\} + o_{p}(\frac{1}{\sqrt{n_{ij}}}) \\ &= \tau_{\varphi} \left\{ \frac{1}{n_{j}} \sum_{m=n_{i}+1}^{n_{ij}} \varphi\left[F(Z_{m})\right] - \frac{1}{n_{i}} \sum_{m=1}^{n_{i}} \varphi\left[F(Z_{m})\right] \right\} + o_{p}(\frac{1}{\sqrt{n}}). \end{aligned}$$

If D_{Δ} is $p \times 1$ vector of $\hat{\Delta}_{ij}s$, $p = \binom{k}{2}$; **R** is $n \times 1$ vector of random variables $\varphi[F(Z_m^{n_i})]$, $i = 1, \ldots, k$ and $m = 1, \ldots, n_i$, then we can write

$$\boldsymbol{D}_{\Delta} = au_{\varphi} \boldsymbol{L} \boldsymbol{R} + o_p(\frac{1}{\sqrt{n}}),$$

where \boldsymbol{L} is a $p \times n$ matrix given by

$$m{L} = egin{bmatrix} -rac{1}{n_1} & rac{1}{n_2} & m{0}_{n_3} & m{0}_{n_4} & \dots & m{0}_{n_{k-1}} & m{0}_{n_k} \ -rac{1}{n_1} & m{0}_{n_2} & rac{1}{n_2} & m{0}_{n_3} & m{0}_{n_4} & \dots & m{0}_{n_{k-1}} & m{0}_{n_k} \ dots & dots &$$

•

Thus, the asymptotic var-cov matrix of \boldsymbol{D}_Δ is

$$\Sigma_{\boldsymbol{D}_{\Delta}} = \tau_{\varphi}^{2} \boldsymbol{L} Var(\boldsymbol{R}) \boldsymbol{L}' = \tau_{\varphi}^{2} \boldsymbol{L} \boldsymbol{L}',$$

where $\Sigma_{\mathbf{R}} = I_n$.

Note that

$$RSM = \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) \hat{\Delta}_{ij} = \sum_{i < j} \sum_{i < j} a_{ij} \hat{\Delta}_{ij} = \mathbf{A}' \mathbf{D}_{\Delta}$$
$$= \tau_{\varphi} \mathbf{A}' \mathbf{L} \mathbf{R} + o_p(\frac{1}{\sqrt{n}}),$$

 $\mathrm{so},$

where A is $p \times 1$ vector of $a_{ij}s$. Proof: Note from above derivations, we have

$$A'L(A'L)' = k^2 \sum_{j=1}^k \frac{a_j^2}{n_j},$$

which is non-negative.

$$\boldsymbol{A}'\boldsymbol{L}_{1\times n} = k \begin{bmatrix} \underline{a_1} & \underline{a_2} & \dots & \underline{a_k} \\ \overline{n_1 n_1} & \overline{n_2 n_2} & \dots & \underline{a_k} \\ n_k n_k \end{bmatrix}.$$

Since $\frac{n_j}{n} \to \lambda_j \in (0,1)$ as $n \to \infty$, n_j and n have the same order, i.e. $O(n_j) = O(n)$. Thus, as $n \to \infty$, every $\frac{a_i}{n_i} \to 0, i = 1, \dots, n$. Therefore,

$$max_{1\leq i\leq n} \left| \boldsymbol{A}' \boldsymbol{L}_i \right| \to 0, as \ n \to \infty.$$

Based on Corollary A.1.1 (Hettmansperger & McKean (2011), Page 447), under H_0

$$RSM = \hat{\tau}_{\varphi} \boldsymbol{A' LR} \xrightarrow{D} N(0, \ \sigma^2_{RSM}),$$

where $\sigma_{RSM}^2 = k^2 \hat{\tau}_{\varphi}^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$.

When $H_0: \theta_1 = \cdots = \theta_k$ is true, then E(RSM) = 0. Hence, under H_0 ,

$$RSM^* = \frac{1}{k\sqrt{\sum_{j=1}^k \frac{a_j^2}{n_j}}} \boldsymbol{A}' \boldsymbol{L} \boldsymbol{R} \xrightarrow{D} Z \sim N(0, 1).$$
(3.10)

The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3) if $RSM^* \ge z_{\alpha}$; otherwise do not reject.

Theorem 3.2. If $H_0: \theta_1 = \cdots = \theta_k$ is true, then E(RSM) = 0. and $Var(RSM) = k^2 \tau_{\varphi}^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$. Then as $n \to \infty$, where $n = \sum_{i=1}^k n_i$,

$$RSM^* = \frac{1}{k\sqrt{\sum_{j=1}^k \frac{a_j^2}{n_j}}} \boldsymbol{A}' \boldsymbol{L} \boldsymbol{R} \xrightarrow{D} Z \sim N(0, 1).$$

3.2.2 Consistency and Asymptotic Power

The null asymptotic distribution of RSM was established in the last section. Thus, a level α test based on RSM for the hypothesis (1.3) coulde be considered.

The *RSM* can be expressed as $\mathbf{A}' \mathbf{D}_{\Delta}$. When H_1 is true, there is a \mathbf{D} which is a vector of the pairwise shifts and $\Delta_{ij} > 0$. Since $O(n_j) = O(n)$, for $\epsilon > 0$ and for large $n, \sigma_{RSM}^2 = k^2 \tau_{\varphi}^2 \sum_{j=1}^k \frac{a_j^2}{n_j} \to 0$, and $0 < z_{\alpha} \sigma_{RSM} < \mathbf{A}' \mathbf{D} - \epsilon$, then the power of the RSM test at fixed alternatives,

$$P_{\boldsymbol{D}}(RSM \ge z_{\alpha}\sigma_{RSM}) \ge P_{\boldsymbol{D}}(\boldsymbol{A}'\boldsymbol{D}_{\Delta} \ge \boldsymbol{A}'\boldsymbol{D} - \boldsymbol{\epsilon}) \ge P_{\boldsymbol{D}}(\left|\boldsymbol{A}'\boldsymbol{D}_{\Delta} - \boldsymbol{A}'\boldsymbol{D}\right| \le \boldsymbol{\epsilon}) \to 1.$$

Thus, RSM is a consistent test.

For a sequence of local alternatives $H_n: \theta_j = \theta_0 + \theta c_j n^{-1/2}$, $(\theta > 0, j = 1, ..., k)$. Under H_n , and F has a density f with $\int f^2(x) dx < \infty$,

$$\Delta_{ij} = (c_j - c_i)\theta n^{-1/2}$$
$$\boldsymbol{D}_n = \boldsymbol{C}\theta n^{-1/2},$$

where C is $p \times 1$ vector of $(c_j - c_i)'s$.

Suppose $D_{\Delta 0}$ is the MWW estimate of D under H_0 , and $D_{\Delta n}$ is the MWW estimate of D under H_n . Since D_{Δ} is translation equivariant, thus,

$$\boldsymbol{D}_{\Delta n} = \boldsymbol{D}_{\Delta 0} + \boldsymbol{D}_n,$$

and

$$\operatorname{Var}(\boldsymbol{D}_{\Delta n}) = \operatorname{Var}(\boldsymbol{D}_{\Delta 0}).$$

Thus, when H_n is true,

$$E(RSM) = E(\boldsymbol{A}'\boldsymbol{D}_{\Delta n}) = \boldsymbol{A}'(\boldsymbol{0} + \boldsymbol{D}_{n}) = \boldsymbol{A}'\boldsymbol{D}_{n}$$

$$\operatorname{Var}(RSM) = \sigma_{RSM}^2.$$

The asymptotic power of the test based on RSM is given by

$$P_{D_n}[RSM^* \ge Z_{\alpha}] = P_{D_n}[RSM \ge Z_{\alpha}\sigma_{RSM}]$$

= $P_{D_n}[\frac{RSM - \mathbf{A}'D_n}{\sigma_{RSM}} \ge Z_{\alpha} - \mathbf{A}'D_n/\sigma_{RSM}]$
= $1 - \Phi(Z_{\alpha} - \theta c_{\varphi 2}),$

where $c_{\varphi 2}$ is the Pitman efficacy of the test. We can write $c_{\varphi 2}$ as

$$c_{\varphi 2} = \mathbf{A}' \mathbf{C} / \sqrt{k \tau_{\varphi}^{2} \sum_{i < j} \sum_{i < j} (a_{j} - a_{i})(\frac{a_{j}}{\lambda_{j}} - \frac{a_{i}}{\lambda_{i}})}$$

$$= \frac{k^{1/2} \sum_{i=1}^{k} a_{i} c_{i}}{\tau_{\varphi} [\sum_{i < j} \sum_{i < j} (a_{j} - a_{i})(\frac{a_{j}}{\lambda_{j}} - \frac{a_{i}}{\lambda_{i}})]^{1/2}}$$

$$= \frac{k^{1/2} \sum_{i=1}^{k} a_{i} c_{i}}{\tau_{\varphi} [k \sum_{i=1}^{k} \frac{a_{i}^{2}}{\lambda_{i}}]^{1/2}}$$

$$= \frac{\sum_{i=1}^{k} a_{i} c_{i}}{[\sum_{i=1}^{k} \frac{a_{i}^{2}}{\lambda_{i}}]^{1/2}} \int f^{2}(x) dx.$$

Since

$$\sum \sum_{i < j} \sum_{i < j} (a_j - a_i) \left(\frac{a_j}{\lambda_j} - \frac{a_i}{\lambda_i}\right) = \sum \sum_{i < j} \left(\frac{a_i^2}{\lambda_i} + \frac{a_j^2}{\lambda_j} - a_i a_j \left(\frac{1}{\lambda_i} + \frac{1}{\lambda_j}\right)\right)$$
$$= (k - 1) \sum_{i=1}^k \frac{a_i^2}{\lambda_i} - \sum_{i=1}^k \frac{a_i}{\lambda_i} (-a_i)$$
$$= \sum_{i=1}^k \frac{a_i^2}{\lambda_i}.$$

Theorem 3.3. If $H_n: \theta_j = \theta_0 + \theta c_j n^{-1/2}, (\theta > 0, j = 1, ..., k)$, and F has a density f with $\int f^2(x) dx < \infty$, then as $n \to \infty$

$$RSM^* \xrightarrow{D} Z \sim N(\theta c_{\varphi}, 1).$$

On the other hand, the Pitman efficacy of the Hettmansperger-Norton test is

$$e = \frac{\sum_{i=1}^{k} a_i (c_i - \bar{c}_w)}{[\sum_{i=1}^{k} \frac{a_i^2}{\lambda_i}]^{1/2}} (12)^{1/2} \int f^2(x) dx$$

$$= \frac{\sum_{i=1}^{k} a_i (c_i - \bar{c}_w)}{[\sum_{i=1}^{k} \frac{a_i^2}{\lambda_i}]^{1/2}} (12)^{1/2} \int f^2(x) dx$$

$$= \frac{\sum_{i=1}^{k} a_i c_i - \sum_{i=1}^{k} a_i \bar{c}_w}{[\sum_{i=1}^{k} \frac{a_i^2}{\lambda_i}]^{1/2}} (12)^{1/2} \int f^2(x) dx$$

$$= \frac{\sum_{i=1}^{k} a_i c_i}{[\sum_{i=1}^{k} \frac{a_i^2}{\lambda_i}]^{1/2}} (12)^{1/2} \int f^2(x) dx = c_{\varphi 2}.$$

Therefore, the efficiency of RSM relative to Hettmansperger-Norton is $c_{\varphi 2}^2/e^2 = 1$, which means that RSM and HN have the same asymptotic local power (Noether, 1955).

3.3 The RSM with Least Square Estimates (RSM_{LS}) Test

The RSM_{LS} test uses the least square estimates $\tilde{\Delta}_{ij} = \bar{Y}_j - \bar{Y}_i$ to estimate the shifts Δ_{ij} . This is our extension of the AT test.

3.3.1 The \mathbf{RSM}_{LS} Test Statistic

The RSM_{LS} test statistic is

$$RSM_{LS} = \sum_{i=1}^{k-1} \sum_{j=i+1}^{k} (a_j - a_i) \tilde{\Delta}_{ij} = \sum_{i < j} \sum_{a_{ij} \tilde{\Delta}_{ij} = \mathbf{A}' \tilde{\mathbf{D}},$$

where the LS estimate of $\Delta_{ij} = \mu_{Y_j} - \mu_{Y_i}$ is $\tilde{\Delta}_{ij} = \bar{Y_j} - \bar{Y_i}$.

$$\begin{split} Cov(\tilde{\Delta}_{ij}, \tilde{\Delta}_{i'j'}) &= Cov(\bar{Y}_j - \bar{Y}_i, \bar{Y}_{j'} - \bar{Y}_{i'}) \\ &= Cov(\bar{Y}_j, \bar{Y}_{j'}) - Cov(\bar{Y}_j, \bar{Y}_{i'}) - Cov(\bar{Y}_i, \bar{Y}_{j'}) + Cov(\bar{Y}_i, \bar{Y}_{i'}) \\ &= I(j = j') \frac{\sigma_j^2}{n_j} - I(j = i') \frac{\sigma_j^2}{n_j} - I(i = j') \frac{\sigma_i^2}{n_i} + I(i = i') \frac{\sigma_i^2}{n_i}. \end{split}$$

Let σ^2 be the common variance of all Y's, then

$$Cov(\tilde{\Delta}_{ij}, \tilde{\Delta}_{i'j'}) = \sigma^2 \left[\frac{I(j=j')}{n_j} - \frac{I(j=i')}{n_j} - \frac{I(i=j')}{n_i} + \frac{I(i=i')}{n_i}\right],$$

Since $n_i, i = 1, ..., k$ has the same order as n, thus as $n \to \infty$

$$\sqrt{n_i}(\bar{Y}_i - \mu_{Y_i}) \xrightarrow{D} N(0, \sigma^2),$$

and $\tilde{\Delta}_{ij}$ is approximately $N(\Delta, \sigma^2(1/n_i + 1/n_j))$.

Let \boldsymbol{V} be a $k\times 1$ vector

$$\boldsymbol{V} = \begin{bmatrix} \sqrt{n_1}((\bar{Y}_1 - \mu_{Y_1})) \\ \vdots \\ \sqrt{n_k}((\bar{Y}_k - \mu_{Y_k})) \end{bmatrix},$$

and if \boldsymbol{L} is a $p \times k$ matrix,

$$\boldsymbol{L} = \begin{bmatrix} -\frac{1}{\sqrt{n_1}} & \frac{1}{\sqrt{n_2}} & 0 & 0 & \dots & 0 & 0 \\ -\frac{1}{\sqrt{n_1}} & 0 & \frac{1}{\sqrt{n_3}} & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & -\frac{1}{\sqrt{n_{k-1}}} & \frac{1}{\sqrt{n_k}} \end{bmatrix},$$

and \boldsymbol{D} is a $p\times 1$ vector

$$m{D} = egin{bmatrix} \mu_{Y_2} - \mu_{Y_1} \ \mu_{Y_3} - \mu_{Y_1} \ dots \ \mu_{Y_k} - \mu_{Y_{k-1}} \end{bmatrix},$$

then $\tilde{D} = LV + D$ and $RSM_{LS} = A'LV + A'D$, where A is $p \times 1$ vector of $a'_{ij}s$. Hence,

$$\boldsymbol{A'L(A'L)'} = k^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$$

which is non-negative.

Since

$$\boldsymbol{A'L}_{1 \times k} = k \begin{bmatrix} \frac{a_1}{\sqrt{n_1}} & \frac{a_2}{\sqrt{n_2}} & \dots & \frac{a_k}{\sqrt{n_k}} \end{bmatrix}$$

and since n_j and n have the same order, as $n \to \infty$, every $a_i/\sqrt{n_i} \to 0, i = 1, \dots, k$. Thus,

$$max_{1 \le i \le n} \left| \boldsymbol{A'} \boldsymbol{L}_i \right| \to 0, as \ n \to \infty.$$

Based on Corollary A.1.1 (HM Page 447)

$$RSM_{LS} = \boldsymbol{A'}\boldsymbol{L}\boldsymbol{V} + \boldsymbol{A'}\boldsymbol{D} \xrightarrow{D} N(\boldsymbol{A'}\boldsymbol{D}, \ \sigma_{RSM_{LS}}^2),$$

where $\sigma_{RSM_{LS}}^2 = k^2 \sigma^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$. When $H_0: \theta_1 = \cdots = \theta_k$ is true, then $E(RSM_{LS}) = 0$. Hence, under H_0 ,

$$RSM_{LS}^* = \frac{1}{k\sigma\sqrt{\sum_{j=1}^k \frac{a_j^2}{n_j}}} (\boldsymbol{A'}\boldsymbol{L}\boldsymbol{V} + \boldsymbol{A'}\boldsymbol{D}) \xrightarrow{D} Z \sim N(0, 1).$$
(3.11)

The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3) if $RSM_{LS}^* \ge z_{\alpha}$; otherwise do not reject.

3.3.2 Consistency and Asymptotic Power

For a level α test based on RSM_{LS} for the hypothesis (1.3), if H_1 is true, all elements in D are greater than 0. Since $O(n_j) = O(n)$, $\sigma_{RSM_{LS}}^2 = k^2 \sigma^2 \sum_{j=1}^k \frac{a_j^2}{n_j} \to 0$ as $n \to \infty$. If n is sufficiently large, we have

$$0 < z_{\alpha} \sigma_{RSM_{LS}} < \boldsymbol{A}' \boldsymbol{D} - \epsilon.$$

Hence,

$$P_{\boldsymbol{D}} [RSM_{LS}^* \ge z_{\alpha}] = P_{\boldsymbol{D}} \left[\boldsymbol{A}' \tilde{\boldsymbol{D}} \ge z_{\alpha} \sigma_{RSM_{LS}} \right]$$
$$\ge P_{\boldsymbol{D}} \left[\boldsymbol{A}' \tilde{\boldsymbol{D}} \ge \boldsymbol{A}' \boldsymbol{D} - \epsilon \right]$$
$$\ge P_{\boldsymbol{D}} \left[\left| \boldsymbol{A}' \tilde{\boldsymbol{D}} - \boldsymbol{A}' \boldsymbol{D} \right| \le \epsilon \right] \to 1,$$

thus the RSM_{LS} test is consistent.

Under H_n : $\theta_j = \theta_0 + \theta c_j n^{-1/2}$, $(\theta > 0, j = 1, ..., k)$, F has a density f with $\int f^2(x) dx < \infty$, so that

$$\Delta_{ij} = (c_j - c_i)\theta n^{-1/2}$$
$$\boldsymbol{D}_n = \boldsymbol{C}\theta n^{-1/2},$$

where C is $p \times 1$ vector of $(c_j - c_i)'s$. Thus under H_n , $RSM_{LS} \xrightarrow{D} N(\mathbf{A}'\mathbf{D}_n, \sigma_{RSM_{LS}}^2)$. The asymptotic power of the test based on RSM_{LS} is given by

$$P_{\boldsymbol{D}_n}[RSM_{LS}^* \ge Z_{\alpha}] = P_{\boldsymbol{D}_n}[RSM_{LS} \ge Z_{\alpha}\sigma_{RSM_{LS}}]$$

$$= P_{\boldsymbol{D}_n}[\frac{RSM_{LS} - \boldsymbol{A}'\boldsymbol{D}_n}{\sigma_{RSM_{LS}}} \ge Z_{\alpha} - \boldsymbol{A}'\boldsymbol{D}_n/\sigma_{RSM_{LS}}]$$

$$= 1 - \Phi(Z_{\alpha} - \theta c_{LS}),$$

where c_{LS} is the Pitman efficacy of the test, and

$$c_{LS} = A'C/\sqrt{k\sigma^{2}\sum_{i

$$= \frac{k^{1/2}\sum_{i=1}^{k}a_{i}c_{i}}{\sigma[\sum_{i

$$= \frac{k^{1/2}\sum_{i=1}^{k}a_{i}c_{i}}{\sigma[k\sum_{i=1}^{k}\frac{a_{i}^{2}}{\lambda_{i}}]^{1/2}}$$

$$= \frac{\sum_{i=1}^{k}a_{i}c_{i}}{\sigma[\sum_{i=1}^{k}\frac{a_{i}^{2}}{\lambda_{i}}]^{1/2}}.$$$$$$

3.4 Comparisons of New Methods

In previous sections we showed that for a fixed alternative, the RAT test, the RAT test, the RAT test, the RSM test and the RSM_{LS} test are all consistent. Thus, the power of these tests tends to one as the sample size increases. In this section, we compare all these methods by using the asymptotic relative efficiency (ARE). Recall that the

Pitman efficacies of four new methods are:

$$c_{RAT} = \frac{\sum h_j c_j}{\sqrt{n} \tau_{\varphi} (\sum \frac{h_j^2}{n_j})^{1/2}}$$

$$c_{RAThn} = \frac{\sum a_j c_j}{\sqrt{n} \tau_{\varphi} (\sum \frac{a_j^2}{n_j})^{1/2}}$$

$$c_{RSM} = \frac{\sum a_j c_j}{\sqrt{n} \tau_{\varphi} (\sum \frac{a_j^2}{n_j})^{1/2}}$$

$$c_{RSMls} = \frac{\sum a_j c_j}{\sqrt{n} \sigma (\sum \frac{a_j^2}{n_j})^{1/2}}.$$

It's obvious that ARE(RAThn, RSM)=1 and ARE(RAThn, RSMls)= $\sigma^2/\tau_{\varphi}^2$. Also ARE(RSM, RSMls)=ARE(RAThn, RSMls). The ARE of the RAT and RAThn tests is

$$ARE(RAT, RAThn) = \frac{(\sum h_j c_j)^2}{n\tau_{\varphi}^2 \sum \frac{h_j^2}{n_j}} / \frac{(\sum a_j c_j)^2}{n\tau_{\varphi}^2 \sum \frac{a_j^2}{n_j}} = \frac{(\sum h_j c_j)^2 \sum \frac{a_j}{n_j}}{\sum \frac{h_j^2}{n_j} (\sum a_j c_j)^2}$$

And for all sample sizes are 5, ARE(RAT,RAThn)=0.9423. Thus, in the following simulation results the RAT test is about 94% efficient as the RAThn test.

3.5 Simulation Results

Ten thousand simulations were run with data which was generated from a normal, right skewed contaminated normal distribution and a log-F distribution to get the empirical powers of all methods, respectively. Eight types of the test, JT (Jonckheere, 1954) and (Terpstra, 1952), SP (McKean et al., 2001), AT (Abelson and Tukey, 1963), HN (Hettmansperger and Norton, 1987), RAT, RAThn (RAThnT is RAThn with t test, RAThnN is RAThn with z test), RSM (RSMT is RSM with t test, RSMN is RSM with z test) and RSMIs (RSMIsT is RSMIs with t test and RSMIsN is RSMIs with N test), were compared in this section.

Four simulations with different situations were run: when errors follow a normal distribution N(0, 1), when errors follow a symmetric heavy tailed contaminated normal distribution CN(0.25, 100), when errors follow a highly right skewed distribution, skewed contaminated normal distribution SCN(0.25, 1, 100), and when errors follow

a heavier-tailed and positively skewed GF(1, 0.1) distribution. New methods were run with Wilcoxon scores under a normal distribution. For the remaining three distributions, new methods were run with Wilcoxon scores, and optimal scores.

3.5.1 With N(0, 1) Underlying Distribution

Table 10 an Figure 13 show the empirical power for all methods with normal errors. The z test of RSMls (RSMlsN) dominates all other methods, and there is no significant difference among JT, SP, HN, RAThnT, RAThnN, RSMT and RSMN. The RAThn test outperforms the RAT test.

Method		Empiric	al Power	
	0	2.11	8.44	18.99
JT	.1007	.5447	.9261	.9972
SP	.0963	.5374	.9255	.9972
AT	.0985	.5365	.9264	.9968
HN	.0994	.5455	.9278	.9973
RAT	.0976	.5141	.9052	.9945
RAThnT	.1007	.5314	.9163	.9960
RAThnN	.1074	.5478	.9230	.9969
RSMT	.1003	.5286	.9180	.9963
RSMN	.1069	.5459	.9250	.9973
RSMlsT	.1007	.5536	.9365	.9982
RSMlsN	.1083	.5714	.9412	.9984

Table 10: Empirical power under a Normal distribution for all methods.

Empirical Power Curves for 10% Tests



Figure 13: Plot of power curves under a Normal distribution for all methods.

3.5.2 With CN(0.25, 100) Underlying Distribution

We ran a CN(0.25, 100) distribution with both Wilcoxon scores and bent4 (.25,.75,-1,1) scores. Table 12 and Figure 15 show that the RAT test and the RAThn test both have higher power than the JT test when using bent4 scores.

Method		Empirical Power										
	0	2.11	8.44	18.99	33.76	52.75	75.96	103.39	135.04			
JT	.1109	.3452	.6233	.8332	.9150	.9579	.9784	.9888	.9946			
SP	.1067	.3357	.6052	.8097	.8914	.9371	.9616	.9752	.9851			
AT	.1037	.1729	.2622	.3845	.4886	.5887	.6736	.7626	.8283			
HN	.1073	.3149	.5753	.7796	.8586	.9149	.9393	.9620	.9722			
RAT	.0730	.2176	.4594	.7091	.8426	.9233	.9588	.9741	.9849			
RAThnT	.0763	.2291	.4728	.7245	.8522	.9275	.9599	.9757	.9867			
RAThnN	.0840	.2437	.4927	.7383	.8615	.9275	.9619	.9777	.9875			
RSMT	.0986	.2551	.4831	.7171	.8356	.9097	.9444	.9654	.9814			
RSMN	.1054	.2681	.5030	.7302	.8440	.9145	.9478	.9675	.9819			
RSMlsT	.1053	.1697	.2582	.3705	.4666	.5731	.6523	.7473	.8041			
RSMlsN	.1135	.1801	.2709	.3859	.4826	.5897	.6644	.7603	.8144			

Table 11: Empirical power under CN(0.25, 100) with Wilcoxon scores.

Empirical Power Curves for 10% Tests (CN-wscores)



Figure 14: Plot of power curves under CN distribution with Wilcoxon scores.

Method		Empirical Power										
	0	2.11	8.44	18.99	33.76	52.75	75.96	103.39	135.04			
JT	.1014	.3421	.6341	.8283	.9225	.9638	.9801	.9882	.9916			
SP	.0969	.3321	.6187	.8046	.8987	.9442	.9632	.9747	.9803			
AT	.1010	.1831	.2724	.3790	.4939	.5917	.6872	.7584	.8146			
HN	.0977	.3201	.5846	.7732	.8661	.9192	.9453	.9606	.9664			
RAT	.1069	.3327	.6302	.8372	.9384	.9733	.9839	.9898	.9918			
RAThnT	.1128	.3474	.6492	.8488	.9394	.9723	.9833	.9896	.9918			
RAThnN	.1208	.3621	.6644	.8585	.9429	.9736	.9835	.9901	.9918			
RSMT	.1640	.3864	.6354	.8100	.9026	.9471	.9681	.9790	.9864			
RSMN	.1718	.3981	.6488	.8177	.9060	.9493	.9649	.9798	.9868			
RSMlsT	.0989	.1763	.2676	.3738	.4757	.5686	.6681	.7387	.8023			
RSMlsN	.1077	.1878	.2797	.3910	.4900	.5821	.6818	.7534	.8123			

Table 12: Empirical power under CN(0.25, 100) with bent4 scores.

Empirical Power Curves for 10% Tests (CN-bent4)



Figure 15: Plot of power curves under CN distribution with bent4 scores.

3.5.3 With SCN(0.25, 1, 100) Underlying Distribution

Table 13 and Figure 16 show that under SCN(0.25,1,100) RAT, RAThn and RSM with Wilcoxon scores outperform AT and RSMls. Among RAT, RAThn and RSM, the z test RAThn (RAThnN) has the highest power.

Method		Empirical Power										
	0	2.11	8.44	18.99	33.76	52.75	75.96	103.39				
JT	.1005	.3297	.6139	.8144	.8976	.9431	.9693	.9789				
SP	.0968	.3195	.5938	.7887	.8697	.9145	.9409	.9555				
AT	.0967	.1519	.2170	.2866	.3737	.4543	.5351	.6127				
HN	.1004	.3263	.6015	.7945	.8738	.9184	.9431	.9569				
RAT	.0686	.1758	.3722	.5963	.7562	.8564	.9158	.9406				
RAThnT	.0768	.1967	.3942	.6161	.7666	.8591	.9140	.9322				
RAThnN	.0814	.2056	.4130	.6320	.7816	.8676	.9192	.9354				
RSMT	.0864	.2106	.4010	.6150	.7610	.8509	.9084	.9303				
RSMN	.0916	.2233	.4187	.6304	.7742	.8592	.9134	.9332				
RSMlsT	.1000	.1562	.2295	.2998	.3831	.4641	.5501	.6224				
RSMlsN	.1084	.1684	.2414	.3136	.3971	.4794	.5653	.6378				

Table 13: Empirical power under SCN(0.25, 1, 100) with Wilcoxon scores.

Empirical Power Curves for 10% Tests (Wilcoxon scores)



Figure 16: Plot of power curves under SCN distribution with Wilcoxon scores.

Table 14 and Figure 17 show the empirical powers of RAT, RAThn and RSM wiht bent1 scores under SCN(0.25,1,100). The RAThn has the highest power of all methods, and has no significant difference with JT.

Method					Empir	ical Pow	ver			
	0	2.11	8.44	18.99	33.76	52.75	75.96	103.39	135.04	170.91
JT	.1043	.3310	.6148	.8018	.9005	.9413	.9632	.9759	.9834	.9893
SP	.1005	.3222	.5989	.7757	.8703	.9146	.9368	.9558	.9624	.9718
AT	.1002	.1555	.2167	.2848	.3656	.4539	.5283	.6078	.6773	.7346
HN	.1057	.3282	.6079	.7817	.8747	.9167	.9385	.9572	.9632	.9736
RAT	.0909	.2599	.5345	.7496	.8777	.9291	.9514	.9686	.9742	.9798
RAThnT	.0973	.2781	.5564	.7633	.8816	.9288	.9505	.9686	.9743	.9820
RAThnN	.1028	.2910	.5718	.7729	.8847	.9327	.9524	.9696	.9755	.9827
RSMT	.1431	.3151	.5614	.7412	.8470	.8997	.9279	.9488	.9571	.9681
RSMN	.1507	.3276	.5742	.7504	.8543	.9035	.9308	.9504	.9577	.9687
RSMlsT	.1016	.1563	.2275	.2962	.3742	.4662	.5441	.6224	.6857	.7527
RSMlsN	.1101	.1677	.2383	.3100	.3882	.4819	.5608	.6364	.7008	.7654

Table 14: Empirical power under SCN(0.25, 1, 100) with bent1 scores.

Empirical Power Curves for 10% Tests (Bent1 scores)



Figure 17: Plot of power curves under SCN distribution with bent1 scores.

3.5.4 With GF(1, .1) Underlying Distribution

If the error distribution is contained in the log-F class, e has a $GF(2m_1, 2m_2)$ distribution. If $(m_1, m_2) = (1, 1)$, then the e has a logistic distribution. Kalbfleisch and Prentice (Kalbfleisch and Prentice, 1980) discussed the class for $m_1, m_2 \ge 1$, Hettmansperger and McKean (Hettmansperger and McKean, 2011) extended the class to m_1 , $m_2 > 0$ for heavier-tailed error distributions. Errors in this section were generated from a heavy-tailed and positively skewed distribution GF(1,.1). Wilcoxon scores, $\log GF(1,.1)$ scores and bent1(.25,-2,1) scores were used in the simulation. In this section, only JT, RAT and RAThn were compared.

Table 15 and Figure 18 show the level and power of JT, RAT, RAThnT and RAThnN with Wilcoxon scores. The power of JT is slightly higher than RAT, RAThnT and RAThnN.

Method		Empirical Power										
	0	0 2.11 8.44 18.99 33.76 52.75 75.96 103.39 135.04										
JT	.1076	.1463	.2064	.2551	.3249	.3969	.4774	.5464	.6186			
RAT	.1025	.1338	.1783	.2186	.2726	.3420	.4033	.4807	.5512			
RAThnT	.1033	.1365	.1848	.2260	.2872	.3581	.4213	.4963	.5636			
RAThnN	.1103	.1444	.1956	.2373	.3020	.3704	.4388	.5142	.5814			

Table 15: Empirical power under GF(1, 0.1) with Wilcoxon scores.





Figure 18: Plot of power curves under GF distribution with Wilcoxon scores.

Table 16 and Figure 19 show the level and power of JT, RAT, RAThnT and RAThnN with loghGF scores. The powers of RAT, RAThnT and RAThnN are higher

than JT, but the empirical α levels are liberial.

Method		Empirical Power											
	0	0 2.11 8.44 18.99 33.76 52.75 75.96 103.39 135.04											
JT	.1046	.1471	.2004	.2596	.3259	.3956	.4692	.5402	.6166				
RAT	.1316	.1897	.2678	.3552	.4520	.5566	.6475	.7359	.8078				
RAThnT	.1365	.1947	.2775	.3687	.4703	.5731	.6627	.7501	.8189				
RAThnN	.1440	.2060	.2910	.3824	.4837	.5868	.6768	.7633	.8289				

Table 16: Empirical power under GF(1, 0.1) with logGF scores.



Empirical Power Curves for 10% Tests (GF-logGF)

Figure 19: Plot of power curves under GF distribution with logGF scores.

Table 17 and Figure 20 show the level and power of JT, RAT, RAThnT and RAThnN with bent1 scores. The power of RAT, RAThnT and RAThnN are higher than JT. RAThnN has the highest power.

Method		Empirical Power											
	0	0 2.11 8.44 18.99 33.76 52.75 75.96 103.39 135.04											
JT	.1053	1053 .1478 .2054 .2555 .3274 .4012 .4637 .5497 .6194											
RAT	.0961	.1504	.2104	.2844	.3811	.4790	.5817	.6770	.7593				
RAThnT	.1006	.1590	.2215	.2928	.3990	.4914	.5990	.6977	.7718				
RAThnN	.1063	.1682	.2324	.3060	.4160	.5099	.6139	.7144	.7862				

Table 17: Empirical power under GF(1, 0.1) with bent1 scores

Empirical Power Curves for 10% Tests (GF-bent1)



Figure 20: Plot of power curves under GF distribution with bent1 scores.

In conclusion, for the parametric-based methods, the RSM_{LS} test is better than the AT test; for the nonparametric-based methods, if the errors follow a right skewed distribution, there is no significant difference among the JT test, the RAT test, the RAThn test and the RSM test, if the errors follow a right skewed and heavy-tailed distribution, the RAT test and the RAThn test are better than the JT test.

3.6 Example

For the example in chapter 1, the new methods test results are given in the following table. RAThn and RSM return robust results, with both p-values less than 0.05. The RSMls result is affected by the outlier, and the p-value is 0.1989, thus it fails to reject the null hypothesis.
1	<u> Table 18:</u>	<u>Test results-ne</u>	$\underline{w} \text{ methods}$
	method	test statistic	p-value
	RAThn	1.8174	0.0446
	RSM	1.8932	0.0389
	RSMls	0.8703	0.1989

CHAPTER 4

MIXED MODELS WITH RANDOM BLOCK EFFECT

In the previous chapters we discussed the methods for an ordered alternative in a one-way linear model. In this chapter we will extend the methods for ordered alternative to a mixed model with two factors. The one factor represents the fixed effect, i.e., the groups, while the second factor is a random block effect. In this case, the observations within each block are dependent random variables. In such designs, the treatment effects have an ordering, such as a increasing intensity of drugs. All the hypothesized ordering information is lost if a Chi-square test or a Friedman test is used. Not considering the order of the hypothesis will cause the loss of efficiency. Page (1963) proposed a test which is based on the weighted rank sums where the ranks are the joint ranking of the observations within blocks. Hettmansperger (1975) proposed a multiple comparison procedure which extended the test statistic of Page and provided the point and interval estimates of the cell location parameters. For more discussion see Hollander (1967), Barlow et al. (1972) and Hollander and Wolfe (1999).

LS estimators and joint-rank estimators are used in the new methods. The method with Abelson-Tukey weights and LS estimators is called Abelson Tukey test with randomized blocks (ATb). The method with Abelson-Tukey weights and jointrank estimators is called JrRAT. If this method contains the block effects in the design matrix, it is called JrRATb. The method with Hettmansperger-Norton weights and joint-rank estimators is called JrRATh. If this method contains the block effects in the design matrix, it is called JrRAThb. Linear mixed-effects model function *lme* was used to get the LS estimates. The estimates of the fixed effects based on the joint ranks (JR) of all residuals in a linear model, with the block correlated continuous error distributions for general score functions is our rank-based fitting procedure. This is the rank-based analog of the traditional LS randomized block design fit. For computation we use the package JRFit (Kloke and McKean, 2014) and (Kloke et al., 2009).

4.1 Simple Mixed Models

Suppose there are k groups and m blocks, where block i has n_i observations; Within block i, let \mathbf{Y}_i , \mathbf{X}_i and \mathbf{e}_i denote the $n_i \times 1$ vector of responses, the $n_i \times p$ design matrix, and the $n_i \times 1$ vector of errors, respectively. Let $\mathbf{1}_{n_i}$ denote the vector of n_i ones. Then the simple mixed model could be written as

$$Y_i = \alpha \mathbf{1}_{n_i} + X_i \beta + \mathbf{1}_{n_i} b_i + \epsilon_i, i = 1, \dots, m,$$
(4.1)

where β is the $k-1 \times 1$ vector of regression coefficients and α is the intercept parameter. The components of ϵ_i are independent and identically distributed and b_i is a continuous random variable which is independent of ϵ_i . Define $e_i = \mathbf{1}_{n_i}b_i + \epsilon_i$. Assume that the random effects b_1, \ldots, b_m are independent and identically distributed.

Note that we can also write the model as

$$\boldsymbol{y_{ij}} = \theta_j + b_i + \epsilon_{ij},$$

where θ_j is the mean (or median) for group j. As in the previous chapters, we are interested in ordered alternatives, so we began with ordered alternatives of the form

$$H_A: \theta_1 \leq \cdots \leq \theta_k$$
, (with at least one strict inequality). (4.2)

4.2 The Page Test

The usual distribution-free test for ordered alternatives is a randomized complete block design known as the Page test. To compute the Page statistic, we first order the k observations within each block. Let r_{ij} denote the rank of y_{ij} in the joint ranking of the observations in the *i*th block and set

$$R_j = \sum_{i=1}^m r_{ij}.$$
 (4.3)

Then the Page statistic L is the weighted rank sums and given by

$$L = \sum_{j=1}^{k} jR_j. \tag{4.4}$$

Notice that this is a Spearman type test statistic.

To test H_0 (1.2) versus the ordered alternative H_1 (1.3) at the α level of significance, the decision rule of Page's test is

Reject H_0 if $L \ge l_{\alpha}$; otherwise do not reject.

The test statistic is distribution free under H_0 . Based on distribution of L under H_0 , approximate values of l_{α} are given in Table A.23 of Hollander & Wolfe (1999). Under H_0 the expected value and the variance of L are

$$E_0(L) = \frac{nk(k+1)^2}{4}$$
$$var_0(L) = \frac{nk^2(k+1)(k^2-1)}{144}$$

The standardized version of L is

$$L^* = \frac{L - E_0(L)}{\sqrt{var_0(L)}}.$$

When H_0 is true, and as n tends to infinity, L^* has an asymptotic N(0, 1). Hence, the asymptotic test is

Reject
$$H_0$$
 if $L^* \ge z_{\alpha}$; otherwise do not reject.

4.3 New Methods

In this section two main types of methods, parametric and nonparametric methods, are introduced. The parametric method is based on the Abelson Tukey test, utilizing LS estimators. We label it the ATb test (Abelson Tukey test with randomized blocks). The robust methods are based on the joint-rank estimators of β in model 4.1; see (Kloke et al., 2009).

The general form of the test statistic of the new method is

$$T^* = \frac{\boldsymbol{w}'\hat{\boldsymbol{\theta}}}{\sqrt{\boldsymbol{w}'var(\hat{\theta})\boldsymbol{w}}},\tag{4.5}$$

where \boldsymbol{w}' is a $k \times 1$ vector of weights (the Abelson Tukey weights and the Hettmansperger-Norton weights are used in this chapter), and $\hat{\boldsymbol{\theta}}$ is the JR estimation of $\boldsymbol{\theta}$, the $k \times 1$ vector of group centers.

To estimate $\hat{\theta}$ we need to fit the linear mixed model

$$Y = X\beta + \mathbf{1}_{n}b + \epsilon. \tag{4.6}$$

where $\boldsymbol{Y} = (\boldsymbol{Y}'_1, \dots, \boldsymbol{Y}'_m)'$, $\boldsymbol{X} = (\boldsymbol{X}'_1, \dots, \boldsymbol{X}'_m)'$, $\boldsymbol{b} = (b_1, \dots, b_m)$ and $\boldsymbol{\epsilon} = (\epsilon_1, \dots, \epsilon_m)$. Let $\hat{\boldsymbol{\beta}}$ denote an estimation of $\boldsymbol{\theta}$ based on a fitting procedure. Let

$$\hat{\boldsymbol{\theta}} = \boldsymbol{E}\hat{\boldsymbol{\beta}} \tag{4.7}$$

where

$$\boldsymbol{E}_{k \times k} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \\ 1 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & 1 \end{bmatrix}.$$
(4.8)

Then a test statistic of the hypothesis (4.2) is given by

$$T^* = \frac{\boldsymbol{w}' \boldsymbol{E} \hat{\boldsymbol{\beta}}}{\sqrt{\boldsymbol{w}' \boldsymbol{E} \boldsymbol{\Sigma}_{\hat{\boldsymbol{\beta}}} \boldsymbol{E}' \boldsymbol{w}}},\tag{4.9}$$

where $\hat{\beta}$ is the estimate of fixed effects, and $\Sigma_{\hat{\beta}}$ is the variance-covariance matrix of the fixed effects.

Based on different weights or different estimators of β , the following methods are generated.

4.3.1 Abelson Tukey Test with Randomized Blocks(ATb)

ATb uses the Abelson-Tukey weights (c) where the estimates of β are computed using the *lme* (Linear Mixed-Effects Models)function in the *nlme* (Linear and Nonlinear Mixed Effects Models) package. The manual on *lme* states that "this function fits a linear mixed-effects model in the formulation described in Laird and Ware (1982) but allowing for nested random effects. The within-group errors are allowed to be correlated and/or have unequal variances." To compute the estimates and the variance-covariance matrix of $\hat{\beta}_{LS}$, the model was fitted by default method, restricted maximum likelihood (REML), which in effect corrects the maximum-likelihood estimator for degrees of freedom. The test statistic is

$$T^* = \frac{c' E \hat{\beta}_{LS}}{\sqrt{c' E \Sigma_{\hat{\beta}_{LS}} E' c}}.$$
(4.10)

For testing H_0 (1.2) versus the ordered alternative H_1 (1.3), large values of T^* lead to the conclusion H_1 . At the α level of significance, the decision rule is

Reject H_0 if $T^* \ge t_{\alpha,n-(k+m-1)}$; otherwise do not reject.

Then two ways were used to compute the p-value

- 1. t test with df = n k m 1
 - n: # of observations
 - m: # of blocks

- k: # of treatment groups
- 1: # of additional parameters estimated in variance-covariance of $\hat{\beta}$
- 2. t test with df = n (k + m 1) [only differs by 2].

4.3.2 Joint Rank Estimator

Joint rank estimators are R estimators based on the joint ranks of all the residuals for fitting linear models with independently distributed errors. Kloke, McKean, and Rashid (2009) extended those estimators to estimating the fixed effects in a linear model with cluster correlated continuous error distributions for general score functions. They called it the JR estimator for joint ranking. For model 4.1, they showed that the asymptotic variance-covariance matrix of $\hat{\beta}_{\varphi}$ is

$$\tau_{\varphi}^{2}(\boldsymbol{X}'\boldsymbol{X})^{-1}\sum_{i=1}^{m}\boldsymbol{X}_{i}'\boldsymbol{\Sigma}_{\varphi,i}\boldsymbol{X}_{i}(\boldsymbol{X}'\boldsymbol{X})^{-1}, \boldsymbol{\Sigma}_{\varphi,i} = (1-\rho_{\varphi})\boldsymbol{I}_{n_{i}} + \rho_{\varphi}\boldsymbol{J}_{n_{i}}, \quad (4.11)$$

where $\rho_{\varphi} = cov \{\varphi[F(e_{11})], \varphi[F(e_{12})]\} = E \{\varphi[F(e_{11})], \varphi[F(e_{12})]\}.$ Let $M = \sum_{i=1}^{m} {n_i \choose 2} - (k-1)$. A simple moment estimator of ρ_{φ} is

$$\hat{\rho}_{\varphi} = M^{-1} \sum_{i=1}^{m} \sum_{l>j} a \left[R(\hat{e}_{il}) \right] a \left[R(\hat{e}_{ij}) \right].$$
(4.12)

Thus an estimate of the asymptotic covariance matrix of the JR estimators is easy to get when plugging in the estimate of τ_{φ} and the estimate of ρ_{φ} (4.11) (Kloke et al., 2009).

Kloke and McKean (Kloke and McKean, 2014) have developed an R package jrfit, which returns the rank-based estimation and inference for mixed models. Like the Rfit package, jrfit provides variaty of score functions, and the default option is Wilcoxon scores. For our computation, we used version 0.03. This package can be found at https://www.biostat.wisc.edu/ kloke/.

4.3.3 Nonparametric Methods

In the following two subsections, all new methods are based on the joint-rank estimates of β and the variance-covariance matrix of β . Two types of the design matrixes are performed in later simulation studies.

1. The design matrix is only composed of the group effect.

$$\boldsymbol{X}_{1,n\times(k-1)} = \begin{bmatrix} \boldsymbol{0} & \boldsymbol{0} & \dots & \boldsymbol{0} \\ \boldsymbol{1}_{n_2} & \boldsymbol{0} & \dots & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{1}_{n_3} & \dots & \boldsymbol{0} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{0} & \boldsymbol{0} & \dots & \boldsymbol{1}_{n_k} \end{bmatrix}.$$
 (4.13)

The centered design matrix based on X_1 , we denote as D_1 .

2. The random blocks effects are also contained in the $(k + m - 1) \times (k + m - 1)$ design matrix given by

$$\boldsymbol{E_2} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & 1 & 0 & \dots & 0 \\ 0 & \dots & \dots & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 1 \end{bmatrix}.$$
(4.14)

 Let

$$\mathbf{X}_{2,n\times(k+m-1)} = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{B} \\ \mathbf{1}_{n_2} & \mathbf{0} & \dots & \mathbf{0} & \mathbf{B} \\ \mathbf{0} & \mathbf{1}_{n_3} & \dots & \mathbf{0} & \mathbf{B} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & \mathbf{1}_{n_k} & \mathbf{B} \end{bmatrix}$$
(4.15)

where \boldsymbol{B} is

$$\boldsymbol{B}_{n_{j}\times(m-1)} = \begin{bmatrix} 0 & 0 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix},$$
(4.16)

with $j = 1, \ldots, k$. Denote by D_2 , the centered design matrix obtained from X_2 .

In the following methods, two procedures are used to estimate the variancecovariance matrix of $\hat{\beta}$; these are sandwich estimation and compound symmetry structure. The sandwich estimation procedure yields asymptotically consistent covariance matrix estimates. In the compound symmetry structure, all the variances are equal and all the covariances are equal. In the *jr fit* function the default variance-covariance matrix is the sandwich estimation.

The following two methods use the Abelson-Tukey weights c along with the joint-rank estimates of β . The test statistic with the different design matrix will be slightly changed:

1. JrRAT

The JrRAT method uses design matrix D_1 , and the test statistic is

$$T_1^* = \frac{\boldsymbol{c}' \boldsymbol{E} \hat{\boldsymbol{\beta}}}{\sqrt{\boldsymbol{c}' \boldsymbol{E} \boldsymbol{\Sigma}_{\hat{\boldsymbol{\beta}}} \boldsymbol{E}' \boldsymbol{c}}}.$$
(4.17)

2. JrRATb

The JrRATb method uses design matrix D_2 , and the test statistic is

$$T_2^* = \frac{\boldsymbol{c}' \boldsymbol{E}_2 \hat{\boldsymbol{\beta}}_b}{\sqrt{\boldsymbol{c}' \boldsymbol{E}_2 \boldsymbol{\Sigma}_{\hat{\boldsymbol{\beta}}_b} \boldsymbol{E}_2' \boldsymbol{c}}}.$$
(4.18)

The subscript b here is for fitting fixed and blocks.

The following two methods use the Hettmansperger-Norton weights along with the joint-rank estimates of β_1 . Note that $\beta = (\alpha, \beta'_1)'$.

3. JrRAThn

The JrRAThn method uses design matrix D_1 , and the test statistic is

$$T_3^* = \frac{k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_1}{k \sqrt{\boldsymbol{a}' \boldsymbol{\Sigma}_{\hat{\boldsymbol{\beta}}_1} \boldsymbol{a}}}.$$
(4.19)

4. JrRAThnb

The JrRAThnb method uses design matrix D_2 , and the test statistic is

$$T_4^* = \frac{k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_{1b}}{k \sqrt{\boldsymbol{a}' \boldsymbol{\Sigma}_{\hat{\boldsymbol{\beta}}_{1b}} \boldsymbol{a}}}.$$
(4.20)

Where $\Sigma_{\hat{\beta}_1} = \Sigma_{\hat{\beta}_1[2:k,2:k]}$. For both types of the design matrixes, the test statistic formulas are the same, but with different estimates for β s and the variance covariance matrices.

Asymptotic theory for these test statistics under the null hypothesis and local alternatives follows as in chapter 3, which uses the asymptotic theory derived in Kloke et al. (2009).

For testing H_0 (1.2) versus the ordered alternative H_1 (1.3), large values of T^* lead to the conclusion H_1 . For each procedure, at the α level of significance, three tests were compired in the simulation study:

- 1. Reject H_0 if $T^* \ge Z_{\alpha}$; otherwise do not reject.
- 2. Reject H_0 if $T^* \ge t_{\alpha,n-k-m-1}$; otherwise do not reject.
- 3. Reject H_0 if $T^* \ge t_{\alpha,n-(k+m-1)}$; otherwise do not reject.

4.4 Simulation Studies

In this section, 10,000 simulations were run with four different data sets. These have different block sizes (m) or treatment groups (k). The m random block effects were randomly generated from $N(0, \sigma_b^2)$; the $n = m \times k$ random errors were randomly generated from $N(0, \sigma_{\epsilon}^2)$. The data set is composed of both the random effects and the random errors, and within each block they have the same block effect. The variance components σ_b^2 and σ_{ϵ}^2 were set to 4 and 1, respectively. The first three data sets have the same block size, 10, but with different treament groups 3, 4 and 7. The fourth data set consists of 15 blocks and 7 treatment groups. Simulation studies on all methods that were mentioned in previous sections will be summarized.

For the simulation studies, the joint-rank estimate of β and the variance-covariance matrix of β with model (4.6) were computed by using the R package *jrfit* (version 0.03).

Table 19 summarizes the empirical alphas and the power of this study with 10 blocks and 3 treatment groups. For both types of design matrix ($D_1 \& D_2$), JrRAT and JrRAThn return exactly the same results for the two types of variance covariance matrix, respectively. Additionally they are not very good on compound symmetry structure but good on sandwich structure. For the first type of design matrix, the methods with a compound symmetry structure variance covariance matrix are slightly liberal. For the second type of design matrix, both methods with both variance covariance matrix work well except the z test. The ATb method works well on both degrees of freedom, and it can significantly outperform the Page method with an empirical power of .4504 and .4531. This is in contrast to .4166 at level .10. JrRATb and JrRAThnb both outperform the Page test. The ATb method has the highest power among all appropriate methods.

Table 20 summarizes the empirical alphas and the power of this study with 10 blocks and 4 treatment groups. For the first type design matrix (D_1), the methods with both compound symmetry structure and sandwich variance covariance matrix are liberal. For the second type design matrix (D_2), z test works well with compound symmetry structure variance covariance matrix; additionally the t tests work well with the sandwich variance covariance matrix. The ATb method works well on both degrees of freedom, and it outperforms the Page method with a power of .4817 and .4828, as opposed to .4417 at level .10. JrRATb and JrRAThnb both outperform the Page test. The ATb method has the highest power among all appropriate methods.

Table 21 summarizes the empirical alphas and the power of this study with 10 blocks and 7 treatment groups. For the first type of design matrix (D_1) , the methods with both compound symmetry structure and sandwich variance covariance matrix are

liberal. For the second type of design matrix (D_2) , the t test works well with the compound symmetry structure variance covariance matrix; all tests with sandwich variance covariance matrix are liberal. The ATb method works well on both degrees of freedom, and it outperforms the Page method with a power of .5650 and .5653, as contrasted with a power of .5382 at level .10. JrRATb and JrRAThnb both outperform the Page test. The JrRAThnb method has the highest power among all appropriate methods.

Table 22 summarizes the empirical alphas and the power of this study with 15 blocks and 7 treatment groups. For the first type of design matrix (D_1), the methods with both compound symmetry structure and sandwich variance covariance matrix are liberal. For the second type of design matrix (D_2), all tests work well with sandwich variance covariance matrix; all tests with compound symmetry structure variance covariance matrix are liberal. The ATb method works well on both degrees of freedom, and it does better than the Page method with a power of .5502 and .5503, as contrasted with a power of .5318 at level .10. The JrRAThnb test outperforms the Page test, and the JrRAThnb method has the highest power among all appropriate methods.

In conclusion, the JrRAThnb test outperforms the Page test among all cases. Among all nonparametric methods the JrRAThnb is the best. The ATb test is a good least square based test.

Method	test	Var-Cov	Df	Empi	rical α		wer
MOMOU	0030	var-00v	DI	10% 5%		1007	50%
	7			10/0	1504	10/0) J/0
	Z		10	.2035	.1524	.5696	.4784
JrRAT	T	CS	16	.1957	.1413	.5555	.4570
	Т		18	.1969	.1425	.5573	.4592
	Z			0963	0533	3081	2740
IrBAT	T	sand	16	.0505	0468	3778	2462
511(111	т	Sand	18	0904	0476	3799	2489
	T		10	.0500	.0470	.0155	.2405
Page				.1143	.0502	.4166	.2475
	Ζ			.1018	.0569	.4176	.2876
JrRATb	Т	CS	16	.0933	.0485	.3999	.2570
	Т		18	.0943	.0493	.4019	.2610
	Z			.1127	.0664	.4342	.3084
JrRATb	Т	sand	16	.1044	.0575	.4149	.2795
	Т		18	.1050	.0585	.4180	.2830
4 (77)	T		10	1015	0510	1501	21.02
ATb	T		10	.1017	.0513	.4504	.3102
ATb	T		18	.1031	.0523	.4531	.3147
	Z			2035	1594	5606	1781
IrP AThn		05	16	1057	1/124	.5090	4570
JII(AI III	T	Co	18	1060	1/125	5573	4502
	T		10	.1909	.1420	.0010	.4092
	Ζ			.0963	.0533	.3981	.2740
JrRAThn	Т	sand	16	.0904	.0468	.3778	.2462
	Т		18	.0908	.0476	.3799	.2489
	Ζ			.1018	.0569	.4176	.2876
JrRAThnb	Т	CS	16	.0933	.0485	.3999	.2570
	Т		18	.0943	.0493	.4019	.2610
	_						
	Z	_		.1127	.0664	.4342	.3084
JrRAThnb	Т	sand	16	.1044	.0575	.4149	.2795
	Т		18	.1050	.0585	.4180	.2830

Table 19: Empirical α s and power(10x3)

Method	test	Var-Cov	Df	Empi	rical α	 Por	wer
			<u> </u>	10% 5%		10%	5%
	7			1724	1181	5624	4502
IrB AT	T	CS	25	1665	.1101	.5624	4335
0110111	Т	65	$\frac{20}{27}$	1669	.1114	.5510	4348
	-		21	.1005	.1110	.0022	. 10 10
	Ζ			.0893	.0476	.3963	.2685
JrRAT	Т	sand	25	.0842	.0422	.3827	.2488
	Т		27	.0845	.0428	.3835	.2502
Page				.1064	.0456	.4417	.2757
	Ζ			.0993	.0505	.4416	.3139
JrRATb	Т	CS	25	.0945	.0438	.4277	.2898
	Т		27	.0949	.0441	.4283	.2915
	_						
	Z			.1089	.0614	.4570	.3326
JrRATb	Т	sand	25	.1034	.0551	.4441	.3108
	Т		27	.1040	.0553	.4448	.3126
ለጥኩ	т		95	1099	0505	1917	990F
ATD	т Т		$\frac{20}{97}$.1036	.0505	.4017	.5505
ALD	T		21	.1044	.0311	.4020	.0099
	Z			1737	1196	5694	4590
JrRAThn	Т	CS	25	.1678	.1123	.5590	.4390
	T		$\overline{27}$.1679	.1127	.5600	.4402
			-				-
	Ζ			.0896	.0473	.3994	.2737
JrRAThn	Т	sand	25	.0842	.0417	.3850	.2538
	Т		27	.0847	.0422	.3858	.2552
	Ζ			.1001	.0515	.4493	.3225
JrRAThnb	Т	CS	25	.0943	.0452	.4373	.2990
	Т		27	.0947	.0458	.4376	.3008
	_						
T D 1	Z		a -	.1098	.0621	.4623	.3396
JrRAThnb	Т	sand	25	.1039	.0561	.4482	.3198
	Έ		27	.1045	.0565	.4488	.3207

Table 20: Empirical α s and power (10x4).

Method	test	Var-Cov	Df	Empi	rical α	Po	wer
				10% 5%		10%	5%
	Ζ			.1406	.0883	.5961	.4760
JrRAT	Т	CS	52	.1384	.0852	.5907	.4659
	Т		54	.1385	.0853	.5908	.4663
	7			0451	0190	9100	1957
	ъ Т	cond	59	.0451	.0169	.3190 .3107	1794
JIIIAI	т Т	Sanu	54	.0435	0170	3107	1785
	T		01	.0400	.0110	.0100	.1100
Page				.1038	.0542	.5382	.3988
	Z			.1040	.0523	.5426	.3965
JrRATb	Т	CS	52	.1014	.0493	.5353	.3858
	Т		54	.1014	.0496	.5356	.3859
	Ζ			.1131	.0653	.5487	.4146
JrRATb	Т	sand	52	.1094	.0613	.5432	.4053
	Т		54	.1094	.0615	.5433	.4059
ATh	Т		52	1026	0519	5650	4151
ATb	T		54	.1026	.0519	.5653	.4153
	Ζ			.1455	.0921	.6263	.5075
JrRAThn	Т	CS	52	.1423	.0888	.6220	.4976
	Т		54	.1423	.0889	.6221	.4982
	7			0440	0109	9419	2019
IrD AThn	Ъ Т	cond	59	.0449 0427	.0192 0174	.0410	.2010 1025
JINAIIII	т Т	sanu	54	.0437	.0174 0175	.3342 3345	1035
	T		94	.0407	.0175	.0040	.1999
	Ζ			.1045	.0558	.5687	.4263
JrRAThnb	Т	CS	52	.1009	.0540	.5620	.4155
	Т		54	.1009	.0540	.5624	.4158
	7			1150	0681	5778	1120
JrB AThnb	T	sand	52	1137	0647	5717	.112 <i>3</i> 4333
9110111110	Т	Sand	54	1137	0647	5719	4336
	Ŧ		01	.1101	.0011	.0110	.1000

Table 21: Empirical α s and power (10x7).

Method	test	Var-Cov	Df	Empi	rical α		wer
	0000	,	21	10% 5%		10%	5%
	7			1991	0703	5667	4496
	Ц Т	00	89	1221	.0703	.5007	.4420 .4257
JIIIAI	т Т	CS	02 84	.1200	.0084	.5020	.4357
	T		04	.1200	.0004	.5020	.4557
	Ζ			.0659	.0301	.4156	.2724
JrRAT	Т	sand	82	.0646	.0293	.4111	.2667
	Т		84	.0646	.0293	.4115	.2670
Page				.1009	.0494	.5318	.3846
	Ζ			.0918	.0460	.5204	.3730
JrRATb	Т	CS	82	.0901	.0439	.5165	.3666
	Т		84	.0902	.0439	.5165	.3668
	Z	_		.1020	.0575	.5262	.3929
JrRATb	Т	sand	82	.0999	.0549	.5232	.3867
	Τ		84	.1001	.0549	.5232	.3867
ለጥኩ	т		00	0069	0450	FEOD	10.19
AID ATL	т Т		02 04	.0908	.0430	.0002	.4048
AID	T		04	.0908	.0431	.0005	.4049
	Z			1285	0716	5962	4747
.IrBAThn	T	CS	82	1260	0694	.5920	4681
01101111111	Ť	65	84	.1268	.0694	.5921	.4683
	Ζ			.0674	.0301	.4450	.2940
JrRAThn	Т	sand	82	.0654	.0287	.4401	.2870
	Т		84	.0655	.0287	.4402	.2872
	Ζ			.0924	.0475	.5481	.4096
JrRAThnb	Т	CS	82	.0915	.0455	.5438	.4035
	Т		84	.0915	.0455	.5440	.4035
	-			1001			11.00
	Z	1	00	.1021	.0550	.5558	.4180
JrRAThnb	Т	sand	82	.1005	.0536	.5513	.4101
	Т		84	.1005	.0536	.5513	.4101

Table 22: Empirical α s and power (15x7).

CHAPTER 5

UNKNOWN PEAK

Consider a dose-response design. In some cases, with increasing dose level, the treatment effect tends to improve, but after some point, advancing dose level tends to mean diminishing performance. The alternative with such a particular pattern of increasing group locations followed by decreasing group locations is called an umbrella alternative. The change point (peak) may be known, but generally it is not known. In such cases the estimate of the peak is of interest. Mack and Wolfe (1981) used a standardized two-sample Mann-Whitney statistic computed between the *i*th group and the remaining k-1 groups combined to estimate the peak. Hettmansperger and Norton (1987) used the index at where the maximum test statistic occurs to estimate the peak, where the maximum is taken over for all p possible patterns of c_1, \ldots, c_k .

Using the methods first introduced in chapter 3, we now explore three methods of estimating the peak. We first describe the methods and then report on the results of a simulation study.

5.1 The Hettmansperger and Norton Method

When the peak is unknown, there are k possible umbrella alternatives, H_A : $\theta_1 \leq \cdots \leq \theta_t \geq \theta_{t+1} \geq \cdots \geq \theta_k$. With equally spaced c_1, \ldots, c_k , where $c_j = j$, for $j = 1, \ldots, t$ and $c_j = 2t - j$ for $j = t + 1, \ldots, k$, there are p possible sets of coefficients c_1, \ldots, c_k . Recall in chapter one that the standardized Hettmansperger and Norton test statistic is

$$V^* = (\frac{12}{n+1})^{1/2} \frac{V}{(\sum a_j^2/\lambda_j)^{1/2}},$$

where $V = \sum_{j} a_j \bar{R}_j$, $\bar{R}_j = \frac{1}{n_j} \sum_{j} R_{ij}$, R_{ij} is the value of Y_{ij} in the combined samples $Y_{11}, \ldots, Y_{kn_k}, \lambda_j = \frac{n_j}{n}$, and $\sum_{j} a_j = 0$.

5.2 New Methods

We take equally spaced c_1, \ldots, c_k , where $c_j = j$, for $j = 1, \ldots, t$ and $c_j = 2t - j$ for $j = t + 1, \ldots, k$. For a given test statistic T_t , we use the index of $\max T_t^*$ to estimate the peak, where the maximum is taken over all possible patterns of c_1, \ldots, c_k and $t = 1, 2, \ldots, p$. We next present three such tests.

5.2.1 RAThn

Recall in chapter 3 that the test statistic of the RAThn is based on the rankbased estimates of the shifts of each group from group 1. Here we use T^* , (6.1), to carry out the test. Thus the T_t^* can be written as

$$T_t^* = \frac{k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_1}{\sqrt{k^2 \tau_{\varphi}^2 \boldsymbol{a}' (\boldsymbol{X}_c' \boldsymbol{X}_c)^{-1} \boldsymbol{a}}},$$
(5.1)

where **a** is a vector of the HN weights $(a_j s)$, and $a_j = \lambda_j (c_j - \sum \lambda_j c_j), j = 1, \dots, k$.

5.2.2 RSM

Recall in chapter 3 the RSM test statistic is based on the MWW estimates of the pairwise shifts. We use the standardized RSM as the T^* . Then the T_t^* can be written as

$$T_t^* = \frac{\sum_{i=1}^{k-1} \sum_{j=i+1}^k (a_j - a_i) \hat{\Delta}_{ij}}{\sqrt{k\tau_{\varphi}^2 \sum \sum_{i < j} (a_j - a_i) (\frac{a_j}{n_j} - \frac{a_i}{n_i})}}.$$
(5.2)

The MWW (or HL) estimate of shift for Δ_{ij} is $\hat{\Delta}_{ij} = med\{Y_{lj} - Y_{l'i}\}, 1 \leq l \leq n_j, 1 \leq l' \leq n_i.$

5.2.3 **RSMls**

Recall in chapter 3 the RSM_{LS} uses the LS estimates of the pairwise shifts. We use the standardized RSM_{LS} as the T^* . Then the T^*_t can be written as

$$T_t^* = \frac{\sum_{i=1}^{k-1} \sum_{j=i+1}^k (a_j - a_i) \hat{\Delta}_{ij}}{\sqrt{k\sigma_{\varphi}^2 \sum \sum_{i < j} (a_j - a_i) (\frac{a_j}{n_j} - \frac{a_i}{n_i})}},$$
(5.3)

where the LS estimate of $\Delta = \mu_{Y_j} - \mu_{Y_i}$ is $\hat{\Delta}_{ij} = \bar{Y}_j - \bar{Y}_i$.

5.3 Simulation Studies

We consider the situation where the number of groups k is 10 and the sample size in each group is 16. Two types of data, a set with a clear peak and a set with a flat peak, were generated both from a normal distribution, and a symmetric heavy tailed contaminated normal distribution.

Initially each group is generated from a normal distribution $N(\mu_i, \sigma)$, for $i = 1, 2, \ldots, 10$. We set $\sigma = 1$. The values of μ_i from the clear peak data set were chosen from 1, 2, ..., 8, 9, 10. For the flat peak data set, the centers of the groups close to the peak should have no significant difference. We set the μ_i s closed to the peak with a 0.5 unit difference from each other. Thus, the μ_i s from the flat peak data set were chosen from 2, 3, ..., 8, 9, 9.5, 10, and the order of the μ_i depends on where the peak is located.

Secondly, each group is generated from a contaminated normal distribution $CN(\epsilon, \mu_i, \sigma^2)$. We set $\epsilon = 0.25$ and $\sigma^2 = 100$. Thus data is generated from mixed $N(\mu_i, 1)$ and $N(\mu_i, 100)$. The values of μ_i are choosen at the same way as the normal distribution case. That is, the values of μ_i from clear peak data set were chosen from 1, 2, ..., 8, 9, 10, and the μ_i s from the flat peak data set were chosen from 2, 3, ..., 8, 9, 9.5, 10.

We consider all ten situations for the peak; i.e., the peak in the *i*th group for *i*. If the peak is located at the 1st group, there is a gradual decrease in the locations of the 10 groups, and the value of μ_i for the respective groups, one through ten, are 10, 9, ..., 1 for the clear peak data set, and are 10, 9.5, 9, ..., 2 for the flat peak

data set. If the peak is located at the 10th group, there is a gradual increase in the locations of the 10 groups, so the value of μ_i are 1, 2, ..., 10 for the clear peak data set and are 2, 3, ..., 9, 9.5, 10 for the flat peak data set.



Figure 21: Plots of clear peak data with different peaks.



Figure 22: Plots of flat peak data with different peaks.

80

If the peak is located at the *i*th group, for $i = 2, \ldots, 9$, the value of μ_i is 10. The value of μ_{i-1} and μ_{i+1} is 9 for the clear peak data set, and is 9.5 for the flat peak data set. The value of μ_{i-2} and μ_{i+2} is 8 for the clear peak data set, and is 9 for the flat peak data set, ..., values are symmetrical on both sides of the peak, and so on. Figure 21 gives sample box plots from the clear peak data set; and figure 22 gives sample box plots from the flat peak data set. All of them are generated from $N(\mu_i, 1)$.

Real	Method	_				# of es	stimates	_	-		10
Peak		1	2	3	4	5	6	7	8	9	10
1	HN RAThn RSM RSM _{ls}	9999 9998 9998 9998 9998	$\begin{array}{c}1\\2\\2\\2\end{array}$								
2	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$	$\begin{array}{c} 31 \\ 6 \\ 6 \\ 4 \end{array}$	9969 9994 9994 9996								
3	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$		7 1	$\begin{array}{c} 9993 \\ 10000 \\ 9999 \\ 10000 \end{array}$							
4	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$				$\begin{array}{c} 10000 \\ 10000 \\ 10000 \\ 10000 \end{array}$						
5	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$					$\begin{array}{c} 10000 \\ 10000 \\ 10000 \\ 10000 \end{array}$					
6	$egin{array}{c} \mathrm{HN} \\ \mathrm{RAThn} \\ \mathrm{RSM} \\ \mathrm{RSM}_{ls} \end{array}$						$\begin{array}{c} 10000 \\ 10000 \\ 10000 \\ 10000 \end{array}$				
7	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$							$\begin{array}{c} 10000 \\ 10000 \\ 10000 \\ 10000 \end{array}$			
8	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$								$\begin{array}{c} 9996 \\ 9999 \\ 9999 \\ 10000 \end{array}$	$\begin{array}{c} 4 \\ 1 \\ 1 \end{array}$	
9	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls} \end{array}$									$9960 \\ 9991 \\ 9990 \\ 9994$	$40 \\ 9 \\ 10 \\ 6$
10	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$									$\begin{array}{c} 1 \\ 1 \\ 2 \end{array}$	$\begin{array}{c} 10000 \\ 9999 \\ 9999 \\ 9999 \\ 9998 \end{array}$

Table 23: Unknown peaks estimation (clear peak $N(\mu_i, 1)$).

Ten thousand simulations were run for each situation under $N(\mu_i, 1)$. In Table 23, for clear peak data, new methods work slightly better than HN. In Table 24, the flat peak simulations, the HN test has an average success rate of 0.9736; the RAThn test has a success rate of 0.9528; the RSM test has an average success rate of 0.9529, and the RSM_{ls} test has an average success rate of 0.9548.

				*		(•		(1 -	, ,,	
Real Peak	Method	1	9	3	4	$\# \text{ of es}_{5}$	timates 6	7	8	0	10
I Cak		1	2	5	4	0	0	'	0	3	10
1	$egin{array}{c} \mathrm{HN} \\ \mathrm{RAThn} \\ \mathrm{RSM} \\ \mathrm{RSM}_{ls} \end{array}$	9075 7851 7847 7896	925 2149 2153 2104								
2	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$	$ \begin{array}{r} 106 \\ 9 \\ 9 \\ 9 \\ 9 \end{array} $	9877 9871 9870 9889	$17 \\ 120 \\ 121 \\ 102$							
3	$egin{array}{c} \mathrm{HN} \\ \mathrm{RAThn} \\ \mathrm{RSM} \\ \mathrm{RSM}_{ls} \end{array}$		$255 \\ 40 \\ 40 \\ 30$	$9745 \\ 9960 \\ 9960 \\ 9970$							
4	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$			$30 \\ 21 \\ 21 \\ 15$	9970 9979 9979 9985						
5	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$				1 1	$\begin{array}{c} 10000 \\ 9999 \\ 10000 \\ 9999 \end{array}$					
6	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$						$\begin{array}{c} 10000 \\ 9997 \\ 9997 \\ 9998 \end{array}$	$3 \\ 3 \\ 2$			
7	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$							$9972 \\ 9979 \\ 9979 \\ 9984$	$28 \\ 21 \\ 21 \\ 16$		
8	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$								9747 9952 9954 9962	$253 \\ 48 \\ 46 \\ 38$	
9	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$								9 112 116 100	$9877 \\9879 \\9876 \\9893$	$\begin{array}{c}114\\9\\8\\7\end{array}$
10	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$									$900 \\ 2191 \\ 2177 \\ 2095$	9100 7809 7823 7905

Table 24: Unknown peaks estimation (flat peak under $N(\mu_i, 1)$)

For the flat peak situation, however, from the peak at the 2nd group to the peak at the 9th group, the average success rate of HN is 0.9899; of RAThn is 0.9952; of RSM is 0.9952, and the average success rate of RSM_{ls} is 0.9960. Both Table 23 and Table 24 show that the closer the peak is to the center, the more accurate the estimate will be. The closer the peak is to the two sides, the higher the error rate will be.

Real	Method		# of estimates								
Peak		1	2	3	4	5	6	7	8	9	10
1	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls} \end{array}$	8419 9794 9779 7675	$1557 \\ 206 \\ 221 \\ 2030$	24 292	3						
2	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls} \end{array}$	$1655 \\ 318 \\ 341 \\ 2490$	$7940 \\9631 \\9600 \\5779$	$405 \\ 51 \\ 59 \\ 1679$	52						
3	$egin{array}{c} \mathrm{HN} \\ \mathrm{RAThn} \\ \mathrm{RSM} \\ \mathrm{RSM}_{ls} \end{array}$	$\begin{array}{c}16\\3\\4\\501\end{array}$	$818 \\ 115 \\ 124 \\ 1583$	$9068 \\ 9873 \\ 9856 \\ 6688$	$98 \\ 9 \\ 16 \\ 1221$	7					
4	$egin{array}{c} \mathrm{HN} \\ \mathrm{RAThn} \\ \mathrm{RSM} \\ \mathrm{RSM}_{ls} \end{array}$	97	$\begin{array}{c}1\\1\\192\end{array}$	$296 \\ 51 \\ 64 \\ 1528$	9648 9945 9932 7170	$56 \\ 3 \\ 3 \\ 1008$	5				
5	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls} \end{array}$	19	15	66	$ \begin{array}{r} 88 \\ 18 \\ 23 \\ 1321 \end{array} $	$9877 \\ 9980 \\ 9974 \\ 7532$	$\begin{array}{r} 35\\2\\3\\1021 \end{array}$	21	2	1	2
6	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls} \end{array}$	3	1	1	27	$37 \\ 4 \\ 6 \\ 1051$	$9862 \\ 9976 \\ 9971 \\ 7522$	$101 \\ 20 \\ 23 \\ 1299$	56	24	16
7	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$					8	$52\\2\\5\\990$	9660 9951 9935 7234	$288 \\ 47 \\ 60 \\ 1512$	148	108
8	HN RAThn RSM RSM _{ls}						4	$118 \\ 11 \\ 15 \\ 1248$	$9018 \\ 9851 \\ 9832 \\ 6528$	$847 \\ 134 \\ 148 \\ 1676$	$\begin{array}{c}17\\4\\5\\544\end{array}$
9	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$							45	$388 \\ 48 \\ 50 \\ 1647$	$8009 \\ 9654 \\ 9632 \\ 5894$	$1603 \\ 298 \\ 318 \\ 2414$
10	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$							3	24 305	$1610 \\ 190 \\ 200 \\ 2083$	8366 9810 9800 7609

Table 25: Unknown peaks estimation (clear peak $CN(0.25, \mu_i, 100)$.)

Ten thousand simulations are run for each situation under $CN(0.25, \mu_i, 100)$. For the clear peak simulations, in Table 25, the HN test has an average success rate of 0.8987; the RAThn has an average success rate of 0.9847; the RSM has an average success rate of 0.9831 and the RSM_{ls} test has an average success rate of 0.6963. Thus, both the RAThn and the RSM test work better than the HN test.

Real	Method		# of estimates								
Peak		1	2	3	4	5	6	7	8	9	10
1	HN RAThn RSM RSM _{ls}	$\begin{array}{c} 6556 \\ 6785 \\ 6781 \\ 5508 \end{array}$	3292 3208 3209 3451	$152 \\ 7 \\ 10 \\ 1009$	32						
2	$\begin{array}{c} \mathrm{HN}\\ \mathrm{RAThn}\\ \mathrm{RSM}\\ \mathrm{RSM}_{ls} \end{array}$	$1713 \\ 365 \\ 387 \\ 2568$	7048 8789 8747 4351	$1239 \\ 846 \\ 866 \\ 2836$	243	2					
3	HN RAThn RSM RSM _{ls}	$113 \\ 17 \\ 22 \\ 1029$	$1637 \\ 532 \\ 554 \\ 1943$	7942 9379 9342 5100	$308 \\ 72 \\ 82 \\ 1892$	36					
4	$\begin{array}{c} \mathrm{HN}\\ \mathrm{RAThn}\\ \mathrm{RSM}\\ \mathrm{RSM}_{ls} \end{array}$	$3 \\ 3 \\ 299$	$7\\1\\1\\462$	$905 \\ 487 \\ 514 \\ 2131$	$8956 \\ 9485 \\ 9453 \\ 5714$	$132 \\ 24 \\ 29 \\ 1360$	32	2			
5	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$	75	52	262	$329 \\ 184 \\ 207 \\ 1924$	$9570 \\ 9792 \\ 9764 \\ 6110$	$101 \\ 24 \\ 29 \\ 1437$	111	14	9	6
6	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$	10	3	5	90	$131 \\ 33 \\ 34 \\ 1418$	$9562 \\ 9794 \\ 9777 \\ 6163$	$307 \\ 173 \\ 189 \\ 1936$	254	68	53
7	$\begin{array}{c} \mathrm{HN}\\ \mathrm{RAThn}\\ \mathrm{RSM}\\ \mathrm{RSM}_{ls} \end{array}$		1		3	26	$120 \\ 16 \\ 21 \\ 1386$	$8967 \\ 9514 \\ 9484 \\ 5789$	$903 \\ 469 \\ 492 \\ 2057$	$9\\1\\3\\490$	1 248
8	$egin{array}{c} \mathrm{HN} \\ \mathrm{RAThn} \\ \mathrm{RSM} \\ \mathrm{RSM}_{ls} \end{array}$					1	37	$262 \\ 77 \\ 86 \\ 1876$	7994 9386 9346 5098	$1631 \\ 522 \\ 550 \\ 1894$	$113 \\ 15 \\ 1894 \\ 1094$
9	$\begin{array}{c} \rm HN\\ \rm RAThn\\ \rm RSM\\ \rm RSM_{ls}\end{array}$						1	$2 \\ 1 \\ 251$	$1275 \\ 923 \\ 942 \\ 2954$	$7023 \\ 8688 \\ 8656 \\ 4320$	$1700 \\ 388 \\ 401 \\ 2474$
10	$\begin{array}{c} \mathrm{HN}\\ \mathrm{RAThn}\\ \mathrm{RSM}\\ \mathrm{RSM}_{ls} \end{array}$							21	$151 \\ 9 \\ 10 \\ 1006$	$3211 \\ 3065 \\ 3096 \\ 3391$	$\begin{array}{c} 6638 \\ 6926 \\ 6894 \\ 5582 \end{array}$

Table 26: Unknown peaks estimation (flat peak $CN(0.25, \mu_i, 100)$)

In Table 26, for the flat peak situation, the HN test has an average success rate of 0.8026; the RAThn test has an average success rate of 0.8854; the RSM test has an average success rate of 0.8824 and the RSM_{ls} test has an average success rate of 0.5374.

In conclusion, under normal distribution, there is no significant difference among all four tests. Under contaminated normal distribution, the RAThn test is the best among all four tests.

CHAPTER 6

ONE-WAY LAYOUT WITH COVARIATES

To reduce large error term variances, researchers may have some covariate measurements along with the treatment measurements. The new method RAThn is based on a rank-based full model, thus it is easy to extend to an ANCOVA model. The full ANCOVA model is fit and the same linear combination of treatments is used as in chapter (3.1). Thus the test automatically adjust for the covariates. In particular, the RAThn test remains consistant.

For the model, we also assume that the slopes for the different groups are homogeneous. If this is not true then the ordering of the treatment effects may depend on the location in the covariate space. For heteroscedastic slopes, a point in covariate space may be selected and an ordering could be tested at that point. This generalizes the rank-based picked-point-analyses developed in Watcharotone and McKean et al. (2015).

An investigation of such a procedure may be done in the future, but, for now, it is beyond the scope of this thesis.

Clearly, tests of homogeneous slopes may be conducted as an initial test. We would recommend the rank-based drop in dispersion test as discussed in Schrader and McKean (1977) and, more recently, in chapter 4 of Kloke and McKean (2014).

Based on the results of this test, one may or may not proceed with an ordering procedure. Such an adaptive procedure may be investigated in the future.

6.1 The Covariance Model

Let Y_{ij} $(i = 1, ..., n_j, j = 1, ..., k)$ be the *i*th response for the *j*th treatment. Let Y be a $n \times 1$ vector of Y_{ij} , where $n = \sum n_j$, let θ be a $k \times 1$ vector of θ_j , and let X_{cov} be a $n \times p$ matrix of covariates, where p is the number of covariates. Then the covariance model is written as

$$Y = W\theta + X_{cov}\gamma + e, \tag{6.1}$$

where \boldsymbol{W} is a $n \times k$ matrix in which the *i*th column has ones corresponding to the *i*th sample and zeros otherwise, $\boldsymbol{\gamma}$ is a $p \times 1$ vector of the slop parameters and \boldsymbol{e} is a $n \times 1$ vector of random errors which are assumed to be *iid* from a continuous distribution.

Then,

$$E(Y_{ij}) = \theta_j + x'_{ij}\gamma.$$

For two treatment groups j and j', if the covariates are the same, we have

$$E(Y_{ij}) - E(Y_{ij'}) = \theta_j - \theta_{j'}.$$

6.2 The AT Test

The AT procedure test statistic for covariates case is presented as

$$T = \frac{\sum_{i=1}^{k} h_i \bar{y}_i}{\sqrt{MS(\sum_{i=1}^{k} \frac{h_i^2}{n_i} + \boldsymbol{d}' \boldsymbol{W_x}^{-1} \boldsymbol{d})}},$$

where h_i s are the maxmin contrast coefficients, \bar{y}_i s are the adjusted group means, MS is the mean square residual within groups from the multiple ANCOVA, W_x is the deviation (i.e., centered) score within group sum of products matrix for the covariates, d is the weighted sample covariate means (see (Huitema, 2011), for details of this test statistic).

6.3 The RAThn Test

As in chapters 3 and 4, the RAThn test for the covariates model is also based on the rank-based model. The estimate of β_1 is obtained from fitting the following model

$$Y = W\theta + X_{cov}\gamma + e$$
$$= \begin{bmatrix} 1 & X_1 & X_{cov} \end{bmatrix} \begin{bmatrix} \alpha \\ \beta_1 \\ \gamma \end{bmatrix} + e$$
$$= X\beta + e.$$

The RAThn test statistic is

$$T = k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_1, \tag{6.2}$$

where $\underline{a} = (a_2, a_3, \ldots a_k)'$ and $\sum_{i=1}^k a_i = 0$.

The *var-cov* matrix of $\boldsymbol{\beta}$ is

$$\boldsymbol{\Sigma}_{\beta} = \begin{bmatrix} \kappa_n & -\tau_{\varphi}^2 \bar{\boldsymbol{X}}' (\boldsymbol{X}' \boldsymbol{X})^{-1} \\ -\tau_{\varphi}^2 \bar{\boldsymbol{X}}' (\boldsymbol{X}' \boldsymbol{X})^{-1} & \tau_{\varphi}^2 (\boldsymbol{X}' \boldsymbol{X})^{-1} \end{bmatrix}, \quad (6.3)$$

where $\kappa_n = n^{-1} \tau_s^2 + \tau_{\varphi}^2 \bar{\boldsymbol{X}}'(\boldsymbol{X}'\boldsymbol{X})^{-1} \bar{\boldsymbol{X}}$ and $\tau_s = (2f(\theta_e))^{-1}$. Let

Let

$$\boldsymbol{H} = \begin{bmatrix} \boldsymbol{0}_{(k-1)\times 1} & \mathbf{I}_{k-1} & \boldsymbol{0}_{(k-1)\times p} \end{bmatrix}.$$
 (6.4)

Thus,

$$var(T) = k^2 \hat{\tau}_{\varphi}^2 \mathbf{a}' \mathbf{H} (\mathbf{X}_c^{*'} \mathbf{X}_c^*)^{-1} \mathbf{H}' a, \qquad (6.5)$$

where

$$X^* = \begin{bmatrix} \boldsymbol{X}_1 & \boldsymbol{X}_{cov} \end{bmatrix}.$$

Based on (2.6) the asymptotically distribution of T is

$$T \dot{\sim} N(k \boldsymbol{a}' \boldsymbol{\beta}_1, k^2 \tau_{\varphi}^2 \boldsymbol{a}' \boldsymbol{H} (\boldsymbol{X}_c^{*'} \boldsymbol{X}_c^*)^{-1} \boldsymbol{H}' \boldsymbol{a}).$$
(6.6)

We next state the null asymptotic distribution of T:

Theorem 6.1. If H_0 : $\theta_1 = \cdots = \theta_k$ is true, then E(T) = 0, and $Var(T) = k^2 \hat{\tau}_{\varphi}^2 a' H(\mathbf{X}_c^{*'} \mathbf{X}_c^*)^{-1} H'a$. Then as $n \to \infty$, where $n = \sum_{i=1}^k n_i$,

$$T^* = \frac{k \boldsymbol{a}' \hat{\boldsymbol{\beta}}_1}{\sqrt{k^2 \hat{\tau}_{\varphi}^2 \boldsymbol{a}' \boldsymbol{H}(\boldsymbol{X}_c^{*'} \boldsymbol{X}_c^*)^{-1} \boldsymbol{H}' \boldsymbol{a}}} \xrightarrow{D} Z \sim N(0, 1).$$

The asymptotic test is

Reject H_0 (1.2) in favor of H_A (1.3) if $T^* \ge z_{\alpha}$; otherwise do not reject.

6.4 Simulation Studies

Ten thousand simulations were run to compare the empirical α s and the empirical power of methods mentioned in previous sections. We set k = 5, p = 3, $n_j = 15$. Three covariates were generated from a multi-variate normal distribution $MVN(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, where $\boldsymbol{\mu}' = (0 \ 0 \ 0)$ and

$$\boldsymbol{\Sigma} = \begin{bmatrix} 1 & 0.4 & 0.3 \\ 0.4 & 1 & 0.3 \\ 0.3 & 0.3 & 1 \end{bmatrix}.$$

By solving the power expression

$$p_0 = P\{F(\delta, k-1, n-k-p) \ge F(1-\alpha, k-1, n-k-p)\},\$$

we obtain the noncentrality parameter δ which is approximately 1.81.

Using the same method as in chapter 2, when setting $\sigma^2 = 1$ and $\mu_1 = 0$, we obtained $d = \sqrt{1.81/150}$.

Under H_0 the response observations are generated from Normal(0, 1), CN(0.25, 100)and SCN(0.25, 1, 100) distributions, respectively. The optimal score functions, Wilcoxon scores, bent4 (0.25,0.75,-1,1) and bent1 (0.75,-2,1) are used for the corresponding distribution.

6.4.1With Normal Underlying Distribution

Table 27 and Figure 23 show the simulation results of the AT, RAThnT and RAThnN tests under normally distributed errors. There are no significant differences among all three tests.

	1									
Method]	Empirical Power								
	0	0.60	2.41	5.43						
AT	.0971	.4960	.8973	.9939						
RAThnT	.0979	.4948	.8954	.9934						
RAThnN	.0997	.4996	.8971	.9935						

Table 27: Empirical α and power under Normal distribution.



Empirical Power Curves for 10% Tests (Normal)

Figure 23: Plot of power curves under Normal distribution.

6.4.2With CN(0.25, 100) Underlying Distribution

Tables 28 and 29 and Figures 24 and 25 all show the simulation results of the AT, RAThnT and RAThnN tests under contaminated normally distributed errors. The scores used for the RAThn procedures are Wilcoxon scores and bent4 (.25,.75,-1,1) scores. Both RAThnT and RAThnN have higher power than the AT test.

Method		Empirical Power								
	0	0 0.60 2.41 5.43 9.65 15.08								
AT	.1019	.1604	.2306	.3187	.4057	.5101				
RAThnT	.0854	.2730	.5405	.7905	.9254	.9762				
RAThnN	.0872	.2764	.5451	.7947	.9272	.9766				

Table 28: Empirical α and power under CN(0.25, 100) distribution with Wilcoxon scores.

Empirical Power Curves for 10% Tests (CN-Wilcoxon scores)



Figure 24: Plot of power curves under CN distribution (Wilcoxon scores).



Empirical Power Curves for 10% Tests (CN-Bent4 scores)

Figure 25: Plot of power curves under CN distribution (bent4 scores).

Method		Empirical Power								
	0	0 0.60 2.41 5.43 9.65 15.0								
AT	.0999	.1564	.2289	.3166	.4143	.5097				
RAThnT	.1041	.3519	.6635	.8828	.9753	.9945				
RAThnN	.1067	.3574	.6676	.8854	.9761	.9945				

Table 29: Empirical α and power under CN(0.25, 100) distribution with bent4 scores.

6.4.3 With SCN(0.25, 1, 100) Underlying Distribution

Tables 30 and 31 and Figures 26 and 27 all show the simulation results of the AT, RAThnT and RAThnN tests under skewed contaminated normal distribution. The scores used for the RAThn procedures are Wilcoxon scores and bent1 (.75,-2,1) scores. Both RAThnT and RAThnN have higher power than the AT test.

Table 30: Empirical α and power under SCN distribution with Wilcoxon scores.

Method	Empirical Power							
	0	0.60	2.41	5.43	9.65	15.08		
AT	.1017	.1588	.2280	.3155	.4215	.5207		
RAThnT	.0888	.2732	.5486	.7925	.9232	.9754		
RAThnN	.0899	.2789	.5538	.7972	.9253	.9763		



Empirical Power Curves for 10% Tests (SCN-Wilcoxon scores)

Figure 26: Plot of power curves under SCN distribution (Wilcoxon scores).

Method	Empirical Power							
	0	0.60	2.41	5.43	9.65	15.08		
AT	.1030	.1396	.1907	.2499	.3084	.3862		
RAThnT	.0779	.2310	.4864	.7340	.8847	.9586		
RAThnN	.0799	.2359	.4908	.7383	.8875	.9594		

Table 31: Empirical α and power under *SCN* distribution with bent1 scores.

Empirical Power Curves for 10% Tests (SCN-Bent1 scores)



Figure 27: Plot of power curves under SCN distribution (bent1 scores).

When errors follow a normal distribution, the RAThn test works as well as the AT test. When errors follow a contaminated normal distribution or a skewed contaminated normal distribution, the RAThn test outperforms the AT test.

CHAPTER 7

CONCLUSIONS AND FUTURE WORK

This chapter summarizes the thesis, and outlines the directions for future work.

7.1 Conclusions

Our investigation consisted of a large simulation study with the number of simulations 10,000, the number of groups set at 5 and with group sizes 5. Error distributions included the normal, Student t with degrees of freedom 10, 8, 5, 3, 2 and 1, and a skewed contaminated normal with mean 1, variance 100 and contaminateion rate 0.25. Besides the Abelson-Tukey (AT), for comparison purposes our study included the robust (wellknown) nonparametric procedures: the Jonckheere-Terpstra (JT), the Spearman (SP), and the Hettmansperger-Norton (HN). These robust tests performed much better for the heavy-tailed error distributions. The JT test is even as efficient as AT on normal errors. The AT test, however, is easily extended to general linear and mixed models.

For a general linear model of the form

$$Y = \alpha + X\beta + e,$$

where \mathbf{Y} is the $n \times 1$ vector of responses, \mathbf{X} is the $n \times (k-1)$ design matrix, and \mathbf{e} is the $n \times 1$ vector of error terms. The least squares estimator minimizes the Euclidean distance between \mathbf{Y} and $\hat{\mathbf{Y}}$, the predicted value of \mathbf{Y} . A rank-based estimator is a different measure of distance which is based on the dispersion function of Jaeckel. The only assumption on the distribution of the errors required is that it is continuous. Our initial investigation of the Abelson-Tukey (AT) procedure, showed that the AT test is not robust for error distribuiton with heavy tails. Thus, the goal of this study is to robustify the AT test. We have developed rank-based procedures for ordered alternative models. These procedures, as the AT procedure, are model based and easily extended to mixed model and ANCOVA cases. Several procedures are proposed and tested in this thesis, i.e. the RAT test, the RAThn test, the RSM test and the RSM_{ls} test. The RAT and RAThn tests are rank-based procedures, the estimates of the shifts in location are obtained from a rank-based regression model as the AT's estimates are obtained from a least squares (LS) fit. Thus, these tests are easily extended to mixed model and ANCOVA cases. The RAThn test uses the AT weights as the AT test. The RAThn test uses the HN weights and the Hodges-Lehmann estimates of the shifts. The RSM_{ls} test uses the HN weights and the least square estimates of the shifts.

Asymptotic variance of four test statistics were derived in the thesis. They are $\tau_{\varphi}^2 \sum_{j=1}^k \frac{h_j^2}{n_j}$, $k^2 \tau_{\varphi}^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$, $k^2 \hat{\tau}_{\varphi}^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$, $k^2 \sigma^2 \sum_{j=1}^k \frac{a_j^2}{n_j}$ of the RAT test, the RAThn test, the RSM test and the RSM_{ls}, respectively. Asymptotic distributions of all test statistics were proposed. The asymptotic power of these tests were also proposed. All four tests are consistent. We showed that the Pitman efficacy of the RAT, RAThn, RSM and RSM_{ls} tests are $n^{-1/2} \frac{\sum h_j c_j}{\tau_{\varphi}(\sum h_j^2/n_j)1/2}$, $n^{-1/2} \frac{\sum a_j c_j}{\tau_{\varphi}(\sum a_j^2/n_j)1/2}$, $n^{-1/2} \frac{\sum a_j c_j}{\tau_{\varphi}(\sum a_j^2/n_j)1/2}$ and $n^{-1/2} \frac{\sum a_j c_j}{\sigma(\sum a_j^2/n_j)1/2}$, respectively. When the number of groups and group size both are 5, ARE(RAT, RAThn)= 0.9423. Thus, under this case, the RAT test is as 94.23% efficient as the RAThn test. The RAThn and the RSM tests have the same Pitman efficacy as the HN test, thus, they have the same asymptotic local power. Thus, asymptotically they are equivalent.

To investigate their small sample properties, simulation studies were obtained on general linear models, mixed models, unknown peak and covariance models, respectively. Normal distribution and some heavy tailed and/or skewed distributions, i.e. contaminated normal distribution, skewed contaminated normal distribution and GF distribution, are used as the error distributions to do the simulations. Ten thousand simulations are run for each situation. For general linear models, simulations were run with different situations: when errors follow a normal distribution N(0, 1), when errors follow a symmetric heavy tailed contaminated normal distribution CN(0.25, 100), when errors follow a highly right skewed distribution, skewed contaminated normal distribution SCN(0.25, 100, 1), and when errors follow a heavier tailed and positively skewed GF(1, 0.1) distribution. We set number of groups 5 and group sizes 5. Under the normal distribution, the RAT, RAThn and RSM tests were run with Wilcoxon scores. For the remaining distributions, they were run with Wilcoxon scores and optimal scores. Under the normal distribution, the RSM_{ls} test dominates all other methods. Under the CN(0.25, 100) distribution, the RAT and RAThn tests both have higher power than the JT test when using bent4 scores. Under the SCN(0.25, 100, 1) distribution, the RAThn test has higher power than all other methods except the JT test, and has no significant difference with the JT test when using bent1 scores. Under the GF(1, 0.1) distribution, the RAT and RAThn

For mixed models, two new methods were generated from the RAT and RAThn tests, but using the JRFit package (Kloke and McKean, 2014) to obtain the estimates of the shifts. Thus, they were called JrRAT and JrRAThn. Two types of the design matrixes, only composed of the group effects or composed of the group effects and the block effects, were used in this chapter. They were called JrRAT, JrRAThn, JrRATb and JrRAThnb, respectively. Two types of the variance-covariance matrixes, the sandwich estimation and the compound symmetry structure, were used in the simulation. Four types of the data sets , 10 blocks with 3, 4 and 7 treatment groups and 15 blocks with 7 groups, were used in the simulation. Ten thousand simulations were run to compare the new methods with the Page test, the standard nonparametric test in this situation. The variance components σ_b^2 and σ_{ϵ}^2 were set to 4 and 1, respectively. The simulation results show that the JrRAThnb test outperforms the Page test in all cases.

For unknown peak problems, three new methods, the RAThn, RSM and RSM_{ls} tests, were compared with the HN test. Two types of data, a set with a clear peak and a set with a flat peak, were generated both from a normal distribution, and a symmetric heavy tailed contaminated normal distribution. The number of groups and the group
size were set as 10 and 16, respectively. All simulation results show that the closer the peak is to the two sides, the higher the error rate will be. And except for the flat peak, under the normal distribution case, the RAThn and RSM tests have higher average success rate than the HN test to estimate the peak.

For covariance models, ten thousand simulations under each case were run to compare the RAThn and the AT tests. We set 5 groups, 3 covariates and 15 observations in each group. Under H_0 the response observations were generated from N(0, 1), CN(0.25, 100) and SCN(0.25, 100. 1), respectively. The simulation results show that when errors follow a normal distribution, the RAThn test works as well as the AT test; when errors follow a contaminated normal distribution or a skewed contaminated normal distribution, the RAThn test outperforms the AT test.

All simulation results show that the RAThn test works well, especially when errors follow a heavy-tailed and/or heavy-tailed skewed distributions.

7.2 Future Work

Although the research presented in this thesis have showed the effectiveness of the RAThn test, it could be further developed in the following three ways.

The RAThn test has an advantage that different optimal score functions correspond to different underlying distributions can be used. Thus, the power of the test can be optimized. The choose of the optimal score functions, however, depends on the underlying error distribution which is unknown in practice. Thus, adding an adaptive procedure will be very useful. Such as the Hogg's adaptive procedure which is discussed in the Hogg, McKean and Craig (2013). We are planning future studies of such schemes.

Another future work of this study is adding an adaptive procedure to detect the homogeneity slopes of the covariance model to decide whether or not to proceed to an ordering procedure. We would recommend the rank-based drop in dispersion test as discussed in Schrader and McKean (1977), and, more recently, in chapter 4 of Kloke and McKean (2014).

The third future work of this study is investigating a procedure for the heteroscedastic slopes. We would recommend the rank-based picked-point-analyses developed in Watcharotone and McKean etc. (2015).

-	Index	
-	Acronym 1	

Abbreviation	Definition
JT	Jonckheere-Terpstra test
SP	Spearman test
AT	Abelson-Tukey test
NH	Hettmansperger-Norton test
RAT	Robust Abelson-Tukey test (with AT weights and Wilconson score)
$\operatorname{RATb1}$	Robust Abelson-Tukey test (with AT weights and bent1 score)
m RATb4	Robust Abelson-Tukey test (with AT weights and bent4 score)
m RAThnT	Robust Abelson-Tukey test (with HN weights and t test)
m RAThnN	Robust Abelson-Tukey test (with HN weights and z test)
RSMT	Robust Shao-Mckean test (with HN weights and t test)
RSMN	Robust Shao-Mckean test (with HN weights and z test)
Page	Page test
JrRAT	Jr estimates with AT weights (design matrix only contains group effects)
JrRATb	Jr estimates with AT weights (design matrix contains group effects and block effects)
ATb	Abelson-Tukey test for mixed models
m JrRAThn	Jr estimates with HN weights (design matrix only contains group effects)
${ m JrRAThnb}$	Jr estimates with HN weights (design matrix contains group effects and block effects)

Bibliography

- Abelson, R. and Tukey, J. (1963). Efficient utilization of non-numerical information in quantitative analysis: General theory and the case of simple order, *The Annals of Mathematical Statistics* 34(4): 1347–1369.
- Alvo, M. (2008). Nonparametric tests of hypotheses for umbrella alternatives, The Canadian Journal of Statistics 36: 143–156.
- Archambault, W., Mack, G. and Wolf, D. (1977). K-sample rank tests using pair-specific scoring functions, *Canad. J. Statist.* 5: 195–207.
- Büning, H. (1999). Adaptive jonckheere-type tests for ordered alternatives, Journal of Applies Statistics 26: 541–551.
- Chang, W., McKean, J., Naranjo, J. and Sheather, S. (1999). High-breakdown rank regression, *Journal of the American tatistical Association* **94**: 205–219.
- Chen, Y. (1991). Notes on the mack-wolfe and chen-wolfe tests for umbrella alternatives, Biometrical Journal, Journal of Mathematical Methods in Biosciences 33: 281–290.
- Chen, Y. and Wolfe, D. (1990). A study of distribution-free tests for umbrella alternatives, Biometrical Journal, Journal of Mathematical Methods in Biosciences 32: 47– 57.
- Chernoff, H. and Savage, I. (1958). Asymptotic normality and efficiency of certain nonparametric test statistics, *Annals of Mathematical Statistics* **29**: 972–994.
- Hettmansperger, T. and McKean, J. (2011). Robust Nonparametric Statistical Methods, Taylor & Francis Group, FL:Boca Ratonl.

- Hettmansperger, T. and Norton, R. (1987). Tests for patterned alternatives in k-sample problems, Journal of the American Statistical Association 82(397): 292–299.
- Hollander, M. and Wolfe, A. D. (1999). Nonparametric Statistical Methods, John Wiley & Sons, Inc., New York.
- Huber, P. (1981). Robust Statistics, John Wiley & Sons, Inc., New York.
- Huitema, B. (2011). Analysis of Covariance and Alternatives: Statistical Methods for Experiments, Quasi-Experiments, and Single-Case Studies, John Wiley & Sons, Inc., New York.
- Jaeckel, L. (1972). Estimating regression coefficients by minimizing the dispersion of the residuals, *The Annals of Mathematical Statistics* 43(5): 1449–1458.
- Jonckheere, A. (1954). A distribution-free k-sample test against ordered alternatives, Biometrika (41): 133–145.
- Jureckova, J. (1971). Nonparametric estimate of regression coefficients, The Annals of Mathematical Statistics 42(4): 1328–1338.
- Kalbfleisch, J. and Prentice, R. (1980). The statistical analysis of failure time data, John Wiley & Sons, Inc., New York.
- Kendall, M. and Stuart, A. (1961). Advanced Theory of Statistics, 2, Hafer Publishing Company, New York.
- Kloke, J. and McKean, J. (2012). Rfit: Rank-based estimation for linear models, The R Journal 4(2): 57–64.
- Kloke, J. and McKean, J. (2014). Nonparametric Statistical Methods Using R, Boca Raton, FL: Chapman-Hall.
- Kloke, J., McKean, J. and Rashid, M. (2009). Rank-based estimation and associated inferences for linear models with cluster correlated errors, *Journal of the American Statistical Association* **104**(485): 384–390.

- Kössler, W. (2006). Some c-sample rank tests of homogeneity against umbrella alternatives with unknown peak, Journal of Statistical Computation and SImulation 76: 57–74.
- Kruskal, W. and Wallis, W. (1952). Use of ranks in one-criterion variance analysis, Journal of American Statistical Association 47: 583–621.
- Mack, G. and Wolfe, D. (1981). k-sample rank tests for umbrella alternatives, Journal of the American Statistical Association 76: 175–181.
- Mann, H. and Whitney, D. (1947). On a test of whether one of two random variables is stochastically larger than the other, Ann. Math. Statist. 18: 50–60.
- McKean, J., Naranjo, J. and Huitema, B. (2001). A robust method for the analysis of experiments with ordered treatment levels, *Psychological Report* **89**(2): 267–273.
- McKean, J., Vidmar, T. and Sievers, G. (1989). A robust two-stage multiple comparison procedure with application to a random drug screen, *Biometrics* **45**(4): 1281–1297.
- Noether, G. (1955). On a theorem of pitman, Annals of Mathematical Statistics **26**: 64–68.
- Odeh, R. (1971). On jonckheere's k-sample test against ordered alternatives, *Techno*metrics 13: 912–918.
- Pan, G. (1996). Distribution-free tests for umbrella alternatives, Communications in Statistics-Theory and Methods 25: 3185–3194.
- Puri, M. (1965). Some distribution-free k-sample rank tests of homogeneity against ordered alternatives, Comm. Pure Appl. Math. 18: 51–63.
- Shi, N. (1988). Rank test statistics for umbrella alternatives, Communications in Statistics. Theory and Methods 17: 2059–2073.
- Spearman, C. (1904). The proof and measurement of association between two things, American Journal of Psychology 15: 72–101.

- Terpstra, J. and Magel, R. (2003). A new nonparametric test for the ordered alternative problem, *Nnparametric Statistics* 15: 289–301.
- Terpstra, T. (1952). The asymptotic normality and consistency of kendall's test against trend, when ties are present, *Indagationes Mathematicae* (14): 327–333.
- Tyron, P. and Hettmansperger, T. (1973). A class of non-parametric tests for homogeneity against ordered alternatives, *Annals of Statistics* 6: 1061–1070.